

Additive Error Guarantees for Weighted Low Rank Approximation

FULL PAPER (Supplementary Material)

Abstract

Low-rank approximation is a classic tool in data analysis, where the goal is to approximate a matrix A with a low-rank matrix L so as to minimize the error $\|A - L\|_F^2$. However in many applications, approximating some entries is more important than others, which leads to the weighted low rank approximation problem. However, the addition of weights makes the low-rank approximation problem intractable. Thus many works have obtained efficient algorithms under additional structural assumptions on the weight matrix (such as low rank, and appropriate block structure). We study a natural greedy algorithm for weighted low rank approximation and develop a simple condition under which it yields bi-criteria approximation up to a small additive factor in the error. The algorithm involves iteratively computing the top singular vector of an appropriately varying matrix, and is thus easy to implement at scale. Our methods also allow us to study the problem of low rank approximation under ℓ_p norm error.

1 Introduction

Matrix low rank approximation is one of the most classic dimension reduction methods in data analysis. The standard least squared error version can also be solved efficiently using the singular value decomposition, and we know how to do this in time comparable to the input sparsity [Clarkson and Woodruff, 2017]. Despite its utility, natural variants of low-rank approximation turn out to be intractable. Weighted low-rank approximation is one well studied example: in many applications, some of the entries of a matrix may be less important to approximate than others (e.g., they might be known to be noisy), and thus we may have a weight associated with each entry. While standard least-squares regression for vectors can incorporate weights directly, the matrix version turns out to be challenging. Formally, the weighted low-rank approximation problem is defined as follows: given $A \in \mathbb{R}^{d \times n}$, a non-negative weight matrix $W \in \mathbb{R}^{d \times n}$ and a parameter k , the goal is to find a rank k matrix L that minimizes $\text{Cost}(L)$,

$$\text{Cost}(L) := \sum_{i,j} W_{ij} \cdot (A_{ij} - L_{ij})^2. \quad (1)$$

The problem and its difficulty were recognized as early as [Young, 1941], and it has been well-studied in the ML literature starting with the work of [Srebro and Jaakkola, 2003]. Unlike the unweighted version (which corresponds to $W = \mathbf{1}_{d \times n}$) low-rank approximation, the problem above is NP-hard in general [Gillis and Glineur, 2011]. Much of the early work such as [Srebro and Jaakkola, 2003, Manton et al., 2003, Eriksson and van den Hengel, 2010] thus developed heuristics for the problem. The first provably efficient algorithms were obtained in the work of [Razenshteyn et al., 2016] (see also references therein for work on matrix completion which is closely related). [Razenshteyn et al., 2016] as well as more recent works [Musco et al., 2020, Ban et al., 2019b] assume

that W has low rank, and develop algorithms that achieve a $(1 + \epsilon)$ (multiplicative) approximation to the optimum cost, while having a running time exponential in the rank of W .

Very recently, [Musco et al., 2020] initiated a study of additive error bounds for weighted low rank approximation. Here the goal is to obtain an L' such that $\text{Cost}(L') \leq \text{OPT} + \epsilon \|A\|_F^2$, where OPT is the optimal cost. Additive error guarantees have been a classic notion in the literature on low rank approximation (starting with the seminal work of [Frieze et al., 2004] on sampling for low rank approximation with additive error). Additive guarantees are realistic in applications where the optimal error is a small yet constant fraction of the total mass (e.g., when a low rank approximation may capture 90% of the Frobenius mass).

So far, our discussion has been restricted to error in the squared norm. However, low rank approximation has also been studied in entrywise ℓ_p norms for $p \neq 2$. While any $p > 0$ ensures that the matrix L approximates A , the choice of p determines how the *non-uniformity* in approximation error is penalized. For example, an ℓ_1 penalty allows some errors $|A_{ij} - L_{ij}|$ to be much larger than others (as long as the total sum is small), while as $p \rightarrow \infty$, higher errors are penalized severely. Thus small values of p are used when some entries can be ignored as outliers (e.g., [Candes and Recht, 2008]), while higher values of p ensure a more uniform approximation.

The works of [Song et al., 2017] and [Ban et al., 2019a] develop sketching based algorithms for ℓ_p norm approximation, particularly for $p \in [1, 2]$. They aim to find low-rank approximations whose objective value is $\leq (1 + \epsilon)$ times the optimum. [Chierichetti et al., 2017] develop approximation guarantees in much more generality, for all $p \geq 1$ (including $p = \infty$). Their result gives a simple $O(k \log n)$ multiplicative approximation to the optimal error.

Goals. Our goal in this paper is to consider weighted low rank approximation with ℓ_2 and ℓ_p error objectives and develop efficient and practical algorithms. We prove the efficacy of the greedy procedure under a novel yet natural assumption and establish additive error guarantees.

1.1 Our results

In all our results, we assume that A is the input matrix, and that W is the non-negative weight matrix which has been re-scaled to satisfy $W_{ij} \in [0, 1]$ for all i, j .

Our first result is to develop a simple greedy algorithm that gives an additive error approximation to weighted low-rank approximation. Unlike prior work, our analysis does not require any explicit assumptions on the weight matrix itself. It works as long as the *target matrix* (the intended low rank approximation) has a Frobenius norm not too large compared to A . Formally, our theorem is the following:

Theorem 1. *Suppose there exists a rank k matrix L that satisfies the two conditions: (a) $\text{Cost}(L) \leq \Gamma$ and (b) $\|L\|_F^2 \leq \Lambda \|A\|_F^2$ for some parameters Λ, Γ . Then for any $\epsilon > 0$, there exists an efficient algorithm that outputs a matrix L' of rank at most $O(k\Lambda/\epsilon^2)$ that satisfies*

$$\text{Cost}(L') \leq \Gamma + \epsilon \|A\|_F^2.$$

Remark. Note that the guarantee is not in terms of the optimal error but in terms of Γ . This is because we could have the optimal matrix L^* having a large value of $\|L^*\|_F$, but there may exist an L with only a slightly larger cost, but a much smaller value of $\|L\|_F^2$.

We also note that in the unweighted case ($W = \mathbf{1}_{d \times n}$), the bound on $\|L\|_F / \|A\|_F$ is automatically satisfied: indeed, the ratio is always ≤ 1 . However in the weighted case, there can be pathological cases where a low-rank approximation has a much higher Frobenius norm than A . As an example,

consider the case of $A = W = \mathbf{I}_n$ ($n \times n$ identity). The matrix $L = \mathbf{1}_{n \times n}$ (all ones) is a rank-one matrix that achieves zero weighted approximation error. However, we have $\|L\|_F^2 / \|A\|_F^2 = n$. Informally, our assumption is equivalent to requiring that even the “unimportant” entries in A are not too different in magnitude from the corresponding entries in L , on average. We believe that this is a reasonable assumption when approximating A by L . Moreover, under this assumption, the theorem requires *no structural assumptions* on W (as in prior work).

Remark. It is natural to ask if a dependence on Λ is *necessary* in general. Showing lower bounds in terms of this parameter is an interesting open direction. However, we note that the known hardness results for matrix completion give an evidence for hardness when A, W are sparse (in this case, $\|L\|_F / \|A\|_F$ is $\Theta(n)$). Specifically, [Hardt et al., 2014] show that for matrix completion, given a matrix A which is the restriction to indices Ω of a rank- k matrix L with entries of magnitude $O(1)$, for any constant c , it is hard to construct a matrix B of rank $r = ck$ such that $\sum_{(i,j) \in \Omega} |A_{ij} - B_{ij}|^2 \leq \epsilon n$. (This is assuming the hardness of an appropriate variant of coloring.) Viewing W as the binary mask matrix corresponding to Ω , this also shows the hardness of weighted low rank approximation. The catch is that the amount of additive error allowed above is quite small; it is $\epsilon \|A\|_F^2$ only when the matrix is sparse.

The algorithm is a greedy procedure that iteratively adds a rank 1 matrix to a decomposition, similar to Frank-Wolfe methods (see, e.g., [Clarkson, 2010]). The crux of the analysis is in showing that in spite of potentially bad choices in the past, there exists an update that can significantly improve the decomposition. A powerful feature of our techniques is that we can extend them to weighted approximation with ℓ_p norm error. We study the entrywise ℓ_p version of the objective in (1), defined as

$$\text{Cost}_p(L) = \sum_{i,j} W_{ij} \cdot |A_{ij} - L_{ij}|^p.$$

Here, additive error will correspond to an ℓ_p analog of the Frobenius norm, $\|X\|_{F_p} := (\sum_{i,j} |X_{ij}|^p)^{1/p}$.

Theorem 2. *Let $p > 2$, and suppose there exists a rank k matrix L that satisfies: (a) $\text{Cost}_p(L) \leq \Gamma$ and (b) $\|L\|_{F_p}^p \leq \Lambda \|A\|_{F_p}^p$ for some parameters Λ, Γ . Then for any $\epsilon > 0$, there exists an efficient algorithm that outputs a matrix L' that satisfies $\text{Cost}_p(L') \leq \Gamma + \epsilon \|A\|_{F_p}^p$. Moreover the rank of L' is at most $O\left(\frac{pk^2\Lambda^{2/p}}{\epsilon^{1+\frac{2}{p}}}\right)$.*

We remark that this does not strictly dominate Theorem 1 because of the additional factor of k . For $p \neq 2$, this extra factor appears because the choice of basis for the target low-rank subspace is important to analyzing the greedy algorithm. As we discuss in Section 3, we need to use a carefully chosen basis for our argument.

Another remark is that our ideas only apply to $p > 2$. When $p < 2$, the maximization problem in each iteration of our current approach turns out to be that of computing the hypercontractive norm of a matrix, which is known to be hard [Barak et al., 2012, Ban et al., 2019a].

Our algorithm follows a similar outline as the one for Theorem 1, but it turns out to be much harder to analyze the improvement. We end up using tools from recent works on ℓ_p regression [Adil et al., 2019, Bubeck et al., 2018]. Moreover, finding a rank-one update in each step requires approximately computing the $p \mapsto 2$ operator norm of an appropriate matrix, which can be done efficiently for $p > 2$ using semidefinite programming, as shown by [Nesterov, 1998, Bhattiprolu et al., 2019].

Finally, as discussed in the introduction, even the unweighted version of low-rank approximation with entrywise ℓ_p error has received a lot of interest, and is known to be challenging. Here, we obtain the following additive approximation.

Theorem 3. *Let $p > 2$, and let OPT_k denote the error of the best rank- k approximation of a given matrix A in the entrywise ℓ_p norm. There exists an efficient (polynomial time) algorithm that outputs an L' of rank $O\left(\frac{pk^2}{\epsilon^{1+\frac{2}{p}}}\right)$ that satisfies the error bound*

$$\|A - L'\|_{F_p}^p \leq \text{OPT}_k + \epsilon \|A\|_{F_p}^p.$$

Unlike the previous theorems, this result is *unconditional*. Indeed, it is a simple consequence of Theorem 2 (see Section 3.4). But to the best of our knowledge, such an additive error approximation for ℓ_p low rank approximation was not known for $p > 2$. Given known hardness results for purely multiplicative approximation, it is interesting to study additive error guarantees (see [Ban et al., 2019a]).

Our algorithm for Theorem 3 can be viewed as extending the familiar *iterative peeling* algorithm for ℓ_2 low-rank approximation to the ℓ_p setting, for $p > 2$. The iterative step is different (now involving a $p \mapsto 2$ norm computation), and we obtain an additive error guarantee. The theorem also complements the sketching-based algorithms for obtaining bi-criteria algorithms for $p \in [1, 2)$ from [Ban et al., 2019a]. Finally, note that when the optimal error OPT_k is very small $\ll \frac{\epsilon}{k \log n} \|A\|_{F_p}^p$, the algorithm of [Chierichetti et al., 2017] has a better guarantee than Theorem 3.

1.2 Notation and overview

All the matrix and vector notations used in the paper will be defined at first use. We begin in Section 2 with the greedy algorithm for the weighted Frobenius error. The framework is then extended to the case of weighted ℓ_p norm error in Section 3. The case of unweighted ℓ_p error (Theorem 3) follows as a corollary and is presented in Section 3.4.

2 Algorithm for squared error

We now present the greedy framework that underlies all of our algorithms.

Outline. Our algorithm proceeds by maintaining a low-rank approximation for A and iteratively adding a rank-1 component that ensures sufficient error reduction. This is done by finding a vector \mathbf{z} and subtracting an appropriate multiple of \mathbf{z} from the *residuals* of each column. The analysis proceeds in a column-by-column fashion, and thus we begin with a few useful lemmas about approximating a single column, and present Algorithm 1 and its analysis in Section 2.1.

Our analysis is similar in spirit to the analysis of the greedy algorithm for column subset selection and sparse coding, [Altschuler et al., 2016, Bhaskara and Tai, 2019], but we need a different view in order to incorporate weights for entries. We begin with a few lemmas about approximating a single column using a collection of vectors. Let $a \in \mathbb{R}^d$ be a vector, and $w \in \mathbb{R}^d$ be weights for the coordinates. Define the function $f_w : \mathbb{R}^d \mapsto \mathbb{R}$ as:

$$f_w(v) = \sum_{i \in [d]} w_i (a_i - v_i)^2, \tag{2}$$

where w_i, a_i, v_i denote the i th entries of the corresponding vectors. Next, suppose that x is a vector (which will be our current approximation for a). Assume that x is “locally optimal” in the sense that increasing or decreasing the magnitude of x does not reduce the value of f_w . Formally, x satisfies $\langle \nabla f_w(x), x \rangle = 0$. The gradient has a simple form in our setting, $\nabla f_w(v) = 2w \circ (a - v)$ (recall that \circ denotes the Hadamard or element-wise product). The following lemma shows how moving along a certain direction improves the value of f_w . First, we define

$$g_w(x, u) = \min_{\eta} f_w(x - \eta u), \quad (3)$$

which is the least possible value of f_w that can be obtained by moving from x along the direction u . (As we can set $\eta = 0$, $g_w(x, u)$ is always $\leq f_w(x)$.)

Lemma 4. *Let a, x, w be defined as above, and let $u \in \mathbb{R}^d$ be a vector such that $|\langle \nabla f_w(x), u \rangle| \geq \gamma$ and $\sum_i w_i u_i^2 \leq 1$. Then we have*

$$g_w(x, u) \leq f_w(x) - \frac{\gamma^2}{4}.$$

Proof. By negating u if necessary, we may assume that $\langle \nabla f_w(x), u \rangle \geq \gamma$. Now, the definition of f_w implies that for any η ,

$$\begin{aligned} f_w(x - \eta u) &= \sum_i w_i (a_i - x_i - \eta u_i)^2 \\ &= \sum_i w_i [(a_i - x_i)^2 - 2\eta(a_i - x_i)u_i + \eta^2 u_i^2] \\ &\leq f_w(x) - \eta \langle w \circ (a - x), u \rangle + \eta^2. \end{aligned}$$

In the last step, we used the assumption that $\sum_i w_i u_i^2 = 1$. Since the middle term is precisely $\langle \nabla f_w(x), u \rangle$, which is $\geq \gamma$ by assumption, we have that $f_w(x - \eta u) \leq f_w(x) - \eta\gamma + \eta^2$. Setting $\eta = \gamma/2$, we obtain the conclusion of the lemma. \square

Next, we show a lemma that is central to our argument. It says that if there is some u such that $f_w(u) < f_w(x)$, and if u can be written as a linear combination of some basis vectors using “small” coefficients, then one of the basis directions can lead to a sufficiently large reduction in the value of f_w . Formally,

Lemma 5. *Let $u_1, u_2, \dots, u_k \in \mathbb{R}^d$ be arbitrary vectors, and suppose $u = \sum_j \alpha_j u_j$, where $\sum_j \alpha_j^2 = B$. Let a, x, w be defined as above, and suppose that $f_w(u) < f_w(x)$. Then*

$$\sum_{j=1}^k |\langle \nabla f_w(x), u_j \rangle|^2 \geq \frac{(f_w(x) - f_w(u))^2}{B}.$$

Proof. We first observe that because of the convexity of f_w (it is a non-negative sum of convex functions), we have that

$$f_w(u) \geq f_w(x) + \langle \nabla f_w(x), u - x \rangle = f_w(x) + \langle \nabla f_w(x), u \rangle.$$

The last equality is because of our assumption that scaling x will not improve f_w . Because $f_w(u) < f_w(x)$, this implies that $|\langle \nabla f_w(x), u \rangle| \geq f_w(x) - f_w(u)$. Now, plugging in $u = \sum_j \alpha_j u_j$ and applying Cauchy-Schwartz, we obtain:

$$\left(\sum_j \alpha_j^2 \right) \left(\sum_j |\langle \nabla f_w(x), u_j \rangle|^2 \right) \geq (f_w(x) - f_w(u))^2.$$

The first term is B by definition, and this completes the proof of the lemma. \square

2.1 Algorithm for weighted approximation

The algorithm proceeds as follows: at time step $t = 0, 1, \dots$, an approximation $\mathbf{x}_j^{(t)}$ is maintained for every column \mathbf{a}_j . Unlike in the single column case above, we now have (potentially) different weight vectors \mathbf{w}_j for each column j . We thus define

$$f_j(v) = \sum_{r \in [d]} w_{j,r} (a_{j,r} - v_r)^2, \quad (4)$$

where $w_{j,r}$ denotes the r th coordinate of \mathbf{w}_j (similarly for \mathbf{a}_j). Since our goal is an additive error approximation, an ideal goal is to bring $f_j(v)$ within $\epsilon \|\mathbf{a}_j\|_2^2$ of the optimal approximation for column j , for all j . (Algorithm 1 gives a full description of the procedure.)

Algorithm 1 Weighted low rank approximation with L_2 error

- 1: **Input:** Matrix $A \in \mathbb{R}^{d \times n}$, error parameter ϵ
 - 2: **Output:** Low-rank approximation $L' \in \mathbb{R}^{d \times n}$ whose columns are spanned by a set of vectors Z , with $|Z| = k' := 8k\Lambda/\epsilon^2$.
 - 3: Initialize $Z = \emptyset$, set $\mathbf{x}_j^{(0)} = 0$ for all j
 - 4: **for** $t = 1, 2, \dots, k'$ **do**
 - 5: Using f_j defined in (4), let $\mathbf{z} \in \mathbb{R}^d$, $\|\mathbf{z}\|_2 = 1$ be the vector that maximizes $\sum_j \langle \nabla f_j(\mathbf{x}_j^{(t-1)}), \mathbf{z} \rangle^2$, and add \mathbf{z} to Z
 - 6: **for** each $j \in [n]$ **do**
 - 7: Compute η that minimizes $f_j(\mathbf{x}_j^{(t-1)} + \eta \mathbf{z})$, and set $\mathbf{x}' = \mathbf{x}_j^{(t-1)} + \eta \mathbf{z}$
 - 8: Compute η that minimizes $f_j(\eta \mathbf{x}')$ and set $\mathbf{x}_j^{(t)} = \eta \mathbf{x}'$
 - 9: **end for**
 - 10: **end for**
 - 11: Return Z and the associated low rank approximation L'
-

Remark. Steps 7 and 8 of the algorithm involve a line search. This is easy in our case because the associated functions of η are univariate quadratics.

We start with some notation concerning the target rank- k solution L (as promised by the statement of Theorem 1). Suppose that $L = UV^T$, where the columns of U are orthonormal, and let $\mathbf{u}_j \in \mathbb{R}^d$, $\mathbf{v}_j \in \mathbb{R}^k$ denote the j th columns of U and V^T respectively. Because of the orthonormal columns in U , we have $\|\mathbf{v}_j\|_2 = \|L_j\|_2$, where L_j is the j th column of L . Our first goal is to obtain a column-wise control on $\|L_j\|_2 / \|\mathbf{a}_j\|_2$. Define the column j to be *good* if $\|L_j\|_2^2 / \|\mathbf{a}_j\|_2^2 \leq \Lambda/\epsilon$ and *bad* otherwise. In what follows, we denote by \mathcal{G} the set of all good columns. The following lemma is easy to see.

Lemma 6. *The total mass of the bad columns of A is small. I.e., $\sum_{j \notin \mathcal{G}} \|\mathbf{a}_j\|^2 \leq \epsilon \|A\|_F^2$.*

Proof. Suppose the contrary, and assume that the inequality fails to hold. By the definition of bad, we have that

$$\sum_{j \notin \mathcal{G}} \|L_j\|^2 > \sum_{j \notin \mathcal{G}} \frac{\Lambda}{\epsilon} \|\mathbf{a}_j\|^2 \geq \Lambda \|A\|_F^2.$$

This contradicts our assumption about the bound on $\|L\|_F^2$ (property (b) in Theorem 1). \square

The lemma allows us to focus on the good columns for most of our analysis. We now introduce the following notation to track the progress of the algorithm.

Notation. We denote

$$\delta_j = \frac{f_j(L_j)}{\|\mathbf{a}_j\|_2^2}, \quad \theta_j^{(t)} = \frac{f_j(\mathbf{x}_j^{(t)})}{\|\mathbf{a}_j\|_2^2}. \quad (5)$$

Thus, informally, our goal is to ensure that $\theta_j^{(t)} \geq \delta_j - \epsilon$ on average. We also study the following weighted averages:

$$\delta^* = \frac{\sum_{j \in \mathcal{G}} \|\mathbf{a}_j\|_2^2 \delta_j}{\|A_{\mathcal{G}}\|_F^2}, \quad \psi^{(t)} = \frac{\sum_{j \in \mathcal{G}} \|\mathbf{a}_j\|_2^2 \theta_j^{(t)}}{\|A_{\mathcal{G}}\|_F^2}, \quad (6)$$

where $A_{\mathcal{G}}$ is the submatrix of A comprising only the good columns. The next lemma shows that if $\psi^{(t)} - \delta^*$ is large, then the $(t+1)$ th iteration makes considerable progress. Formally,

Lemma 7. *Suppose that after the t 'th iteration of the algorithm we have $\psi^{(t)} > \delta^*$. Then there exists a \mathbf{z} such that*

$$\sum_{j \in \mathcal{G}} |\langle \nabla f_j(\mathbf{x}_j^{(t)}), \mathbf{z} \rangle|^2 \geq \frac{\epsilon \|A_{\mathcal{G}}\|_F^2 (\psi^{(t)} - \delta^*)^2}{k\Lambda}$$

Proof. The idea will be to prove that one of the $\{\mathbf{u}_i\}_{i \in [k]}$ satisfies the condition of the lemma. We do this by applying Lemma 5 to each of the good columns. Step 8 ensures that the current representation for each column cannot be improved by rescaling, which is essential for applying Lemma 5. Consider any $j \in \mathcal{G}$. This implies that L_j can be written as $\sum_{i \in [k]} \alpha_i \mathbf{u}_i$, where $\sum_i \alpha_i^2 \leq \frac{\Lambda}{\epsilon} \|\mathbf{a}_j\|_2^2$ (indeed the α_i are precisely the entries of the column \mathbf{v}_j). Thus, by applying Lemma 5, we get

$$\begin{aligned} \sum_{i \in [k]} |\langle \nabla f_j(\mathbf{x}_j^{(t)}), \mathbf{u}_i \rangle|^2 &\geq \frac{\epsilon (f_j(\mathbf{x}_j^{(t)}) - f_j(L_j))_+^2}{\Lambda \|\mathbf{a}_j\|_2^2} \\ &= \frac{\epsilon \|\mathbf{a}_j\|_2^2 (\theta_j^{(t)} - \delta_j)_+^2}{\Lambda}. \end{aligned} \quad (7)$$

We first show that the sum of the RHS above over $j \in \mathcal{G}$ is large. By viewing $\frac{\|\mathbf{a}_j\|_2^2}{\|A_{\mathcal{G}}\|_F^2}$ as a probability distribution over the indices $j \in \mathcal{G}$ and using the fact that $\mathbb{E}[X^2] \geq \mathbb{E}[X]^2$, we get

$$\sum_{j \in \mathcal{G}} \frac{\|\mathbf{a}_j\|_2^2}{\|A_{\mathcal{G}}\|_F^2} (\theta_j^{(t)} - \delta_j)_+^2 \geq \left(\sum_{j \in \mathcal{G}} \frac{\|\mathbf{a}_j\|_2^2}{\|A_{\mathcal{G}}\|_F^2} (\theta_j^{(t)} - \delta_j)_+ \right)^2$$

Using the observation that for any real numbers c, d , $(c)_+ + (d)_+ \geq (c + d)_+$ (and generalizing this to a sum of multiple terms), the RHS above can be simplified (using (6)) as

$$\sum_{j \in \mathcal{G}} \frac{\|\mathbf{a}_j\|_2^2}{\|A_{\mathcal{G}}\|_F^2} (\theta_j^{(t)} - \delta_j)_+ \geq (\psi^{(t)} - \delta^*)_+.$$

The RHS is positive by assumption, and thus plugging the above back into (7), we get:

$$\sum_{i \in [k]} \sum_{j \in \mathcal{G}} |\langle \nabla f_j(\mathbf{x}_j^{(t)}), \mathbf{u}_i \rangle|^2 \geq \frac{\epsilon \|A_{\mathcal{G}}\|_F^2 (\psi^{(t)} - \delta^*)^2}{\Lambda}.$$

Thus, by averaging, there exists an index i that satisfies the conclusion of the lemma. This completes the proof. \square

The next lemma bounds the progress after t steps of the algorithm.

Lemma 8. *Let $\epsilon < 1/2$ be a given error parameter. The number of iterations needed to achieve $\psi^{(t)} - \delta^* \leq 2\epsilon$ is $O\left(\frac{k\Lambda}{\epsilon^2}\right)$.*

Proof. Recall that $\psi^{(t)}$ and δ^* only involve the good columns. Define $\beta_t := \psi^{(t)} - \delta^*$, and note that β_t clearly only reduces as t increases. We are done if $\beta_t \leq 2\epsilon$, and thus consider some $t \leq \frac{8k\Lambda}{\epsilon^2}$ and assume that $\beta_t > 2\epsilon$.

We claim that in the next $O(k\Lambda/\epsilon\beta_t)$ steps, the value of β_t reduces by a factor 2. To see this, suppose the contrary.

Now in each iteration, the algorithm finds some \mathbf{z} with $\|\mathbf{z}\| = 1$ such that the total leftover mass (over *all* the columns) reduces by at least the bound given by Lemma 7. This is because the algorithm finds \mathbf{z} that maximizes $\sum_j \langle \nabla f_j(\mathbf{x}_j^{(t-1)}), \mathbf{z} \rangle^2$, and by Lemma 4, this also quantifies the total mass reduction. (Note that we have used the fact that all the weights are $\in [0, 1]$ when applying the Lemma.) Thus, since $\beta_{t'} \geq \beta_t/2$ for all the time steps t' we are considering, the mass reduction is at least

$$\frac{\epsilon \|A_{\mathcal{G}}\|_F^2 \beta_t^2}{4k\Lambda} \geq \frac{\epsilon \|A\|_F^2 \beta_t^2}{8k\Lambda},$$

where we used $\epsilon < 1/2$ and Lemma 6. Thus if this continues for $8k\Lambda/\epsilon\beta_t$ steps, the total mass reduction (which includes the reduction on bad columns) is $\geq \beta_t \|A\|_F^2$. But since $\beta_t > 2\epsilon$ and at most $\epsilon \|A\|_F^2$ of the mass is on the bad columns, this contradicts our assumption that β_t did not reduce by a factor 2.

Thus, we have argued that as long as $\beta_t > 2\epsilon$, it takes $\leq 8k\Lambda/\epsilon\beta_t$ steps for β_t to reduce to $\beta_t/2$. Since $\beta_0 \leq 1$, we have that it takes $\leq 2^j \cdot \frac{8k\Lambda}{\epsilon}$ steps for β_t to reduce from 2^{-j} to $2^{-(j+1)}$. Thus, as the geometric series converges to twice the last term, we have that β_t reduces to $\leq 2\epsilon$ after $\frac{2}{2\epsilon} \cdot \frac{8k\Lambda}{\epsilon}$ steps, completing the proof of the lemma. \square

We can now complete the proof of Theorem 1.

Proof of Theorem 1. Lemma 8 gives us that after $O\left(\frac{k\Lambda}{\epsilon^2}\right)$ steps, we have $\psi^{(t)} - \delta^* \leq 2\epsilon$. Combined with Lemma 6, we have that the overall error in approximation is at most $2\epsilon \|A_{\mathcal{G}}\|_F^2 + \epsilon \|A\|_F^2 \leq 3\epsilon \|A\|_F^2$. This completes the proof (after replacing ϵ by $\epsilon/3$ throughout). \square

3 Low rank approximation with ℓ_p error

The high level outline of our algorithm is similar to the ℓ_2 setting. However, we need the right target decomposition, and need to set up the analysis carefully so as to make the rank-one update at every step efficient.

3.1 Identifying a target decomposition

Let A be the matrix to be approximated and W the weight matrix as before. We make the same assumption: $L = UV^T$ is the target decomposition, and we have $\|L\|_{F_p}^p / \|A\|_{F_p}^p \leq \Lambda$, for some parameter Λ .

Recall that the starting point in our analysis in the case of ℓ_2 error was to decompose L as UV^T using the SVD, so that we have a U with orthonormal columns, and a V such that $\|L_j\| = \|\mathbf{v}_j\|$. Implicit here is the fact that the ℓ_2 norm is rotation invariant (using a different basis U maintains the norm property). Unfortunately, this is not true in the case of ℓ_p norms. A priori, it is not clear if there exists a good decomposition that allows a property such as the above for all the columns, nor is it clear what normalization one should choose for the columns of U . E.g., should they have $\|\cdot\|_p = 1$, or a different norm such as ℓ_2 or the dual of ℓ_p ?

So our first step is to describe the target decomposition and its properties.

Lemma 9. *Let $L \in \mathbb{R}^{d \times n}$ be any rank k matrix with $k \leq \min\{d, n\}$. Then there exists a decomposition $L = UV^T$ into $(d \times k)$ and $(k \times n)$ matrices such that (a) the columns of U satisfy $\|\mathbf{u}_i\|_p = 1$ for all $i \in [k]$, (b) for all $j \in [n]$, the columns of V^T satisfy $\|\mathbf{v}_j\|_\infty \leq \|L_j\|_p$.*

Proof. The proof uses the following simple observation about rank k matrices.

Observation. Let $M \in \mathbb{R}^{d \times n}$ be a rank k matrix. Then there exist a subset S of k columns of M with the property that all the other columns can be expressed as $\sum_{i \in S} \alpha_i M_i$, with $|\alpha_i| \leq 1$ for all i .

The observation follows by an extremal argument, considering the k columns such that the volume of the associated parallelpiped is maximized. We refer the reader to [Chierichetti et al., 2017] (Lemma 2) for a proof. (The argument itself is classic, and the property above is related to the notion of an Auerbach basis in Banach spaces. See [Taylor, 1947, Martini et al., 2001]. One of the early applications of this idea in the CS literature was in the work of [Awerbuch and Kleinberg, 2004].)

For our lemma, we apply the observation above to the matrix M whose columns are $M_j = \frac{L_j}{\|L_j\|_p}$. Let the chosen columns of M be denoted by the vectors \mathbf{u}_i , for $i \in [k]$. Then we have that all the other M_j can be expressed as $\sum_i \alpha_i \mathbf{u}_i$ with $|\alpha_i| \leq 1$, and thus the corresponding L_j can be expressed using coefficients $|\alpha_i| \leq \|L_j\|_p$. By construction, the \mathbf{u}_i have $\|\cdot\|_p = 1$, which completes the proof of the lemma. \square

The lemma allows us to use the framework from Section 2 to develop an iterative algorithm.

3.2 Single vector analysis

The first main step is to obtain analogs of Lemmas 4 and 5

Let $w, a \in \mathbb{R}^d$, and define the function $f_{w,p} : \mathbb{R}^d \mapsto \mathbb{R}$ as:

$$f_{w,p}(v) = \sum_{i \in [d]} w_i |a_i - v_i|^p, \quad (8)$$

where w_i, a_i are the i th entries as before. Observe that the function $f_{w,p}$ is convex (as it is a sum of convex functions). The key to our proof is an appropriate smoothness property for f , which we prove using the following lemma from [Adil et al., 2019].

Lemma 10 (Lemma 4.5 of [Adil et al., 2019]). *Let $y \in \mathbb{R}$ and Δ be any parameter. Then we have*

$$|y + \Delta|^p \leq |y|^p + g\Delta + 2^p \gamma_p(|y|, \Delta),$$

where g is the derivative of $|y|^p$, i.e., $g = p|y|^{p-2}y$, and γ_p is the function (originally introduced in [Bubeck et al., 2018]):

$$\gamma_p(t, \Delta) = \begin{cases} \frac{p}{2} t^{p-2} \Delta^2 & \text{if } |\Delta| \leq t, \\ |\Delta|^p + \left(\frac{p}{2} - 1\right) t^p & \text{otherwise.} \end{cases}$$

Using this lemma, we will be able to show the following analog of Lemma 4.

Lemma 11. *Let a, x, w be defined as above, and let $u \in \mathbb{R}^d$ be a vector such that $|\langle \nabla f_{w,p}(x), u \rangle| \geq \gamma \geq 0$ and $\sum_i w_i |u_i|^p \leq 1$. Then there exists η such that for some constant $c_p = O_p(1)$,*

$$f_{w,p}(x - \eta u) \leq f_{w,p}(x) - \frac{\gamma^2}{c_p (f_{w,p}(x))^{\frac{p-2}{p}}}.$$

Proof. Let η be a parameter that we will choose appropriate. We start by observing that $f_{w,p}(x - \eta u) = \sum_i w_i |a_i - x_i - \eta u_i|^p$. In what follows, we define $y = (a - x)$ for simplicity. Thus, using Lemma 10, we have

$$f_{w,p}(x - \eta u) \leq \sum_i w_i [|y_i|^p - \eta g_i u_i + 2^p \gamma_p(|y_i|, \eta u_i)],$$

where g_i is the gradient $p|y_i|^{p-2}y_i u_i$. We will also use the following upper bound on the function γ_p :

$$\gamma_p(t, \Delta) \leq |\Delta|^p + \frac{p}{2} t^{p-2} \Delta^2. \quad (9)$$

This follows from a simple case analysis from the definition in Lemma 10. By replacing u with $-u$ if necessary, we may assume that the hypothesis $|\langle \nabla f_{w,p}(x), u \rangle| \geq \gamma$ implies that $\sum_i w_i g_i u_i \geq \gamma$. Thus the *decrease* in the value of $f_{w,p}$ is at least

$$D := \eta \gamma - 2^p \sum_i w_i \gamma_p(|y_i|, \eta u_i).$$

The rest of the proof will aim to choose an $\eta > 0$ and show a lower bound on D . Using (9), we have

$$D \geq \eta \gamma - 2^p \sum_i w_i \left(\eta^p |u_i|^p + \frac{p}{2} |y_i|^{p-2} \eta^2 |u_i|^2 \right).$$

The first of the two terms in γ_p is easy to handle, by noting that $\sum_i w_i \eta^p |u_i|^p \leq \eta^p$ (using the hypothesis on u). Thus, let us focus on the other term. We claim that

$$\sum_i w_i |y_i|^{p-2} u_i^2 \leq \left(\sum_i w_i |u_i|^p \right)^{2/p} \left(\sum_i w_i |y_i|^p \right)^{\frac{p-2}{p}}. \quad (10)$$

This follows from Hölder's inequality $\langle \alpha, \beta \rangle \leq \|\alpha\|_\rho \|\beta\|_{\rho'}$, applied to the vectors α, β whose coordinates are

$$|\alpha_i| = w_i^{\frac{2}{p}} |u_i|^2, \quad |\beta_i| = w_i^{\frac{p-2}{p}} |y_i|^{p-2},$$

and $\rho = p/2$ (and the dual norm $\rho' = p/(p-2)$). Now, the first term on the RHS of (10) is bounded by 1 as before. The second term is related to the value of $f_{w,p}$, as by definition, $f_{w,p}(x) = \sum_i w_i |y_i|^p$. Let us write $F = f_{w,p}(x)$ in what follows.

Putting the above observations together, we have that the decrease D satisfies

$$D \geq \eta\gamma - 2^p \left(\eta^p + \frac{p\eta^2}{2} F^{\frac{p-2}{p}} \right) = \eta\gamma - 2^p \eta^p - p2^{p-1} F^{\frac{p-2}{p}} \eta^2.$$

Our choice of η will ensure that the second term is upper bounded by the third term. This is equivalent to $2\eta^{p-2} \leq pF^{(p-2)/p}$. As $p \geq 2$, this will hold as long as $\eta \leq F^{1/p}$.

The value of η we consider is

$$\eta = \frac{\gamma}{p2^{p+1} F^{(p-2)/p}}.$$

Showing that this is $\leq F^{1/p}$ is equivalent to showing that $\gamma \leq p2^{p+1} F^{(p-1)/p}$. By assumption, the gradient term $|\langle \nabla f_{w,p}(x), u \rangle| \geq \gamma$, thus

$$\begin{aligned} \gamma &\leq p \sum_i w_i |y_i|^{p-2} y_i u_i \\ &\leq p \left(\sum_i w_i |y_i|^p \right)^{\frac{p-1}{p}} \left(\sum_i w_i |u_i|^p \right)^{\frac{1}{p}}. \end{aligned}$$

Since the second term is ≤ 1 , we have that $\eta < F^{1/p}$, as desired.

Thus, for the above value of η , we have

$$D \geq \eta\gamma - p2^p F^{\frac{p-2}{p}} \eta^2 \geq \frac{\gamma^2}{p2^{p+2} F^{\frac{p-2}{p}}}.$$

Plugging in the definition of F completes the proof of the lemma. \square

Our analysis will also need an analog of Lemma 5 where f_w is replaced by $f_{w,p}$. This is immediate because the proof only relies on the convexity of f_w , and thus also applies to $f_{w,p}$.

3.3 Algorithm and analysis

Similar to the ℓ_2 case, we define

$$f_{j,p}(v) = \sum_{r \in [d]} w_{j,r} |a_{j,r} - v_r|^p, \quad (11)$$

where w_j is the weight vector for the j th column and $w_{j,r}$ denotes the r th coordinate of w_j (similarly for a_j).

Algorithm. The algorithm for the ℓ_p error case is precisely the same as before, but instead of working with the functions f_j , we work with $f_{j,p}$ (when taking gradients). The main change is

in Step 5 of the algorithm, where instead of finding a vector \mathbf{z} that maximizes $\sum_j \langle \nabla f_j(\mathbf{x}_j^{(t)}), \mathbf{z} \rangle^2$ subject to $\|\mathbf{z}\| = 1$ (which reduces to finding the top singular vector of an appropriate matrix), we now need to solve the following:

$$\max \sum_j \frac{\langle \nabla f_{j,p}(\mathbf{x}_j^{(t)}), \mathbf{z} \rangle^2}{(f_{j,p}(\mathbf{x}_j^{(t)}))^{\frac{p-2}{p}}} \text{ subject to } \|\mathbf{z}\|_p = 1. \quad (12)$$

This can be re-written as finding a vector \mathbf{z} that maximizes $\|M\mathbf{z}\|_2^2$ subject to $\|\mathbf{z}\|_p = 1$, for an appropriate matrix M (which we can construct since we know $\mathbf{x}_j^{(t)}$ and f). This is exactly the problem of computing the so-called $p \mapsto 2$ operator norm of the matrix M . The classic result of [Nesterov, 1998] shows that the problem admits a constant factor approximation. More recently, the work [Bhattiprolu et al., 2019] obtains nearly tight factors for the problem. Both these algorithms are based on a semidefinite programming relaxation for approximating the operator norm, and crucially rely on $p \geq 2$ in their analysis. We summarize these results as follows.

Theorem 12. [Nesterov, 1998, Bhattiprolu et al., 2019] *For any $p \geq 2$, there exists an efficient (polynomial time) algorithm for approximating the $p \mapsto 2$ operator norm of a matrix M to a factor only depending on p (which indeed turns out to be $O(\sqrt{p})$ using the result of [Steinberg, 2005]) Specifically, the algorithm outputs a \mathbf{z} with $\|\mathbf{z}\|_p = 1$, such that the objective value in (12) is $\Omega(\frac{1}{p})$ times the optimum.*

Our analysis once again involves quantities δ_j and $\theta_j^{(t)}$, defined as follows:

$$\delta_j = \frac{f_{j,p}(L_j)}{\|\mathbf{a}_j\|_p^p}, \quad \theta_j^{(t)} = \frac{f_{j,p}(\mathbf{x}_j^{(t)})}{\|\mathbf{a}_j\|_p^p}. \quad (13)$$

We also define weighted averages as before:

$$\delta^* = \frac{\sum_{j \in \mathcal{G}} \|\mathbf{a}_j\|_p^p \delta_j}{\|A_{\mathcal{G}}\|_{F_p}^p}, \quad \psi^{(t)} = \frac{\sum_{j \in \mathcal{G}} \|\mathbf{a}_j\|_p^p \theta_j^{(t)}}{\|A_{\mathcal{G}}\|_{F_p}^p}. \quad (14)$$

The following lemma shows that as long as $\psi^{(t)} - \delta^*$ is large enough, the algorithm makes significant progress.

Lemma 13. *Suppose that after the t 'th iteration of the algorithm we have $\psi^{(t)} > \delta^*$. Then there exists a unit vector \mathbf{z} such that*

$$\sum_{j \in \mathcal{G}} \frac{|\langle \nabla f_{j,p}(\mathbf{x}_j^{(t)}), \mathbf{z} \rangle|^2}{(f_{j,p}(\mathbf{x}_j^{(t)}))^{\frac{p-2}{p}}} \geq \frac{\epsilon^{2/p} \|A_{\mathcal{G}}\|_{F_p}^p (\psi^{(t)} - \delta^*)^2}{k^2 \Lambda^{2/p}}.$$

Proof. The proof follows the structure of that of Lemma 7, and will show that one of the \mathbf{u}_i satisfy the conclusion of the lemma. Consider some good column j .

Our updates ensure that we can apply Lemma 5 (where f_j is replaced by $f_{j,p}$). The value of Λ that we use in the lemma statement is the following: every coefficient used is $\leq \|L_j\|_p$ in magnitude, from Lemma 9. Since j is a good column, this is at most $(\frac{\Lambda}{\epsilon})^{1/p} \|\mathbf{a}_j\|_p$. As there are k terms, the

sum of squared coefficients is bounded by $k \|\mathbf{a}_j\|_p^2 (\frac{\Lambda}{\epsilon})^{2/p}$. Plugging this in, and writing $C = (\frac{\Lambda}{\epsilon})^{2/p}$ for convenience, we obtain:

$$\begin{aligned} \sum_{i \in [k]} |\langle \nabla f_{j,p}(\mathbf{x}_j^{(t)}), \mathbf{u}_i \rangle|^2 &\geq \frac{(f_{j,p}(\mathbf{x}_j^{(t)}) - f_{j,p}(L_j))^2_+}{kC \|\mathbf{a}_j\|_p^2} \\ &= \frac{\|\mathbf{a}_j\|_p^{2p} (\theta_j^{(t)} - \delta_j)_+^2}{kC \|\mathbf{a}_j\|_p^2}. \end{aligned} \quad (15)$$

Thus, since $f_{j,p}(\mathbf{x}_j^{(t)}) = \theta_j^{(t)} \|\mathbf{a}_j\|_p^p$ by definition, we have (after plugging in above and simplifying the exponent of $\|\mathbf{a}_j\|_p$)

$$\begin{aligned} \sum_{i \in [k]} \frac{|\langle \nabla f_{j,p}(\mathbf{x}_j^{(t)}), \mathbf{u}_i \rangle|^2}{(f_{j,p}(\mathbf{x}_j^{(t)}))^{\frac{p-2}{p}}} &\geq \frac{\|\mathbf{a}_j\|_p^p (\theta_j^{(t)} - \delta_j)_+^2}{kC (\theta_j^{(t)})^{\frac{p-2}{p}}} \\ &\geq \frac{\|\mathbf{a}_j\|_p^p (\theta_j^{(t)} - \delta_j)_+^2}{kC}. \end{aligned}$$

The second inequality uses the fact that $p \geq 2$ and $\theta_j^{(t)} \in (0, 1]$. Then, we can sum over the columns $j \in \mathcal{G}$, and mimicking the idea from the proof of Lemma 7 (this time using $\|\mathbf{a}_j\|_p^p / \|A_{\mathcal{G}}\|_{F_p}^p$ as the distribution), we get

$$\sum_{i \in [k]} \sum_{j \in \mathcal{G}} \frac{|\langle \nabla f_{j,p}(\mathbf{x}_j^{(t)}), \mathbf{u}_i \rangle|^2}{(f_{j,p}(\mathbf{x}_j^{(t)}))^{\frac{p-2}{p}}} \geq \frac{\|A_{\mathcal{G}}\|_{F_p}^p (\psi^{(t)} - \delta^*)^2}{kC}.$$

Thus by averaging and plugging in the value of C , one of the \mathbf{u}_i must satisfy the conclusion of the lemma. \square

We will first establish an analog of Lemma 8 for the current setting.

Lemma 14. *Let $\epsilon < 1/2$ be a given error parameter. The number of iterations needed to achieve $\psi^{(t)} - \delta^* \leq 2\epsilon$ is $O\left(\frac{pk^2\Lambda^{2/p}}{\epsilon^{1+2/p}}\right)$.*

Proof. Similar to the ℓ_2 case, define $\beta_t := \psi^{(t)} - \delta^*$, and note that β_t clearly only reduces as t increases. We are done if $\beta_t \leq 2\epsilon$, and thus consider some $t \leq \frac{8pk^2\Lambda^{2/p}}{\epsilon^{2/p}}$ and assume that $\beta_t > 2\epsilon$.

We claim that in the next $O(pk^2\Lambda^{2/p}/\epsilon^{2/p}\beta_t)$ steps, the value of β_t reduces by a factor 2. To see this, suppose the contrary.

In each iteration, the algorithm finds some \mathbf{z} with $\|\mathbf{z}\|_p = 1$ such that the total leftover mass (over all the columns) reduces by at least the bound given by Lemma 7. This is because the algorithm finds \mathbf{z} that is an approximation for the the problem of maximizing $\sum_j \frac{\langle \nabla f_{j,p}(\mathbf{x}_j^{(t)}), \mathbf{z} \rangle^2}{(f_{j,p}(\mathbf{x}_j^{(t)}))^{\frac{p-2}{p}}}$ which we can see is $\Omega(1/p)$ approximation to the optimum by Theorem 12. By Lemma 11, this also quantifies the total mass reduction. (Note that we have used the fact that all the weights are $\in [0, 1]$ when

applying the Lemma.) Thus, since $\beta_{t'} \geq \beta_t/2$ for all the time steps t' we are considering, the mass reduction is at least

$$\frac{\epsilon^{2/p} \|A_{\mathcal{G}}\|_{F_p}^p \beta_t^2}{4pk^2\Lambda^{2/p}} \geq \frac{\epsilon^{2/p} \|A\|_{F_p}^p \beta_t^2}{8pk^2\Lambda^{2/p}},$$

where we used $\epsilon < 1/2$ and the definition of *good* columns. Thus if this continues for $(8pk^2\Lambda^{2/p})/(\epsilon^{2/p}\beta_t)$ steps, the total mass reduction (which includes the reduction on bad columns) is $\geq \beta_t \|A\|_F^2$. But since $\beta_t > 2\epsilon$ and at most $\epsilon \|A\|_F^2$ of the mass is on the bad columns, this contradicts our assumption that β_t did not reduce by a factor 2.

Thus, we have argued that as long as $\beta_t > 2\epsilon$, it takes $\leq 8pk^2\Lambda^{2/p}/\epsilon^{2/p}\beta_t$ steps for β_t to reduce to $\beta_t/2$. Since $\beta_0 \leq 1$, we have that it takes $\leq 2^j \cdot \frac{8pk^2\Lambda^{2/p}}{\epsilon^{2/p}}$ steps for β_t to reduce from 2^{-j} to $2^{-(j+1)}$. Thus, as the geometric series converges to twice the last term, we have that β_t reduces to $\leq 2\epsilon$ after $\frac{2}{2\epsilon} \cdot \frac{8pk^2\Lambda^{2/p}}{\epsilon^{2/p}}$ steps, completing the proof of the lemma. \square

Given this lemma, the proof of Theorem 2 follows as before.

Proof of Theorem 2. First, we can see that from Lemma 14 the number of steps needed to reach $\psi^{(t)} - \delta^* \leq 2\epsilon$ is $O\left(\frac{pk^2\Lambda^{2/p}}{\epsilon^{1+2/p}}\right)$.

Finally, observing that the bound on the total mass of the bad columns carries over to the ℓ_p case, the theorem follows. \square

3.4 Unconditional result for uniform weights

We now show how to deduce Theorem 3 using Theorem 2.

Proof of Theorem 3. We only need to check that the matrix achieving the optimal error (say L^*) satisfies the conditions of Theorem 2. This is true because

$$\|L^*\|_{F_p} = \|(L^* - A) + A\|_{F_p} \leq \|L^* - A\|_{F_p} + \|A\|_{F_p}.$$

By definition, $\|L^* - A\|_{F_p} = \text{OPT}_k$, which is $\leq \|A\|_{F_p}$. This implies that the assumption holds with $\Lambda = 2^p$. \square

4 Experiments

In this section we evaluate our algorithm (*wlra-iter*) for weighted low rank approximation by comparing its performance with three baselines: (a) applying SVD to the matrix A (*svd*) (b) applying SVD to weighted matrix $W \circ A$ (*wsvd*) (c) regularized weighted low rank approximation algorithm with sketching in [Ban et al., 2019b] (*rwlra-sk*). In (c), we use the alternating minimization based algorithm provided in [Ban et al., 2019b]. We present experiments on both synthetic and real data below.

4.1 Synthetic datasets

We conduct two sets of experiments. In the first set, we vary the output rank k' and show how the error changes for each algorithm. In the second set, we demonstrate how the error in each algorithm changes as the signal to noise ratio (SNR) varies: the signal is a low rank matrix and we add Gaussian noise to it. In each experiment, we measure the scaled error $(\sum_{ij} W_{ij}(A_{ij} - Z_{ij})^2) / \|A\|_F^2$ where Z is the solution output by each algorithm (we note that in the experiments in [Ban et al., 2019b], the objective value is plotted instead of the error thus our experiments are not comparable); we average results over 10 independent runs.

We first generate 500×5 dimensional matrices M_1, M_2 with random orthonormal vectors as columns and a diagonal matrix S with diagonal elements $[1, 0.9, (0.9)^2, (0.9)^3, (0.9)^4]$ (normalized). Thus $M = M_1 S M_2^T$ is a rank 5 matrix with $\|M\|_F = 1$. In each experiment we create matrix A by adding a noise matrix N with $N_{ij} \sim \mathcal{N}(0, \sigma^2)$ to M . We set the sketch size parameter in *rwlr-sk* to 100 in all experiments. We generate weight matrices of 500×500 dimension with the following configurations.

- W_1 : Each element is sampled from $\{1, 0.1, 0.01\}$ with probabilities $\{0.85, 0.1, 0.05\}$.
- W_2 : Each element is sampled from $\{1, 0.1, 0.01\}$ with probabilities $\{0.05, 0.1, 0.85\}$.
- W_3 : Each element is sampled from the interval $[0, 1]$ uniformly at random.
- W_4 : Each element is sampled from $\{0, 1\}$ with probabilities 0.3, 0.7.
- W_5 : Elements corresponding to largest 50000 $|A_{ij}|$ s are set to 0, and 1 elsewhere.
- W_6 : Block diagonal is set to 0 where the block size is 100×100 , and 1 elsewhere.
- W_7 : A random binary matrix is first chosen by setting each entry to 1 with probability 0.1 and 0 otherwise. Following this, the first 100 columns of first 150 rows are set to 1.

In the first set of experiments, we plot the error of each algorithm with output rank k' in the list (5, 10, 15, 20, 25, 30, 35, 40, 50, 60). Here we fix $\sigma = 0.005$ (thus $\text{SNR} \approx 0.16$) and $\lambda = 0.05$ for weight matrix settings W_1, W_4, W_5, W_6 and $\lambda = 0.01$ for weight matrix settings W_2, W_3, W_7 in *rwlr-wk*. Figure 1 shows the error rates of each algorithm for different weight matrices.

In the second set of experiments, we plot the error of each algorithm as the SNR is increased from 0.0004 to 4. Here we fix $k' = 50$ and $\lambda = 0.005$ for weight matrix settings W_1, W_2, W_3 and $\lambda = 0.01$ for weight matrix settings W_4, W_5, W_6, W_7 in *rwlr-wk*. We control SNR by changing σ appropriately. Figure 2 shows the error rates of each algorithm for different weight matrices. The results show the greedy procedure achieving small recovery error even in low SNR regimes.

4.2 Real datasets

In this section we compare the performance of *wlr-iter* with *svd* and *wsvd*. We do not include *rwlr-sk* in this set of experiments as it is difficult to tune the parameter λ in *rwlr-sk* and it is not in the scope of this paper. We use following four datasets in this set of experiments.

1. NIPS Conference Papers 1987-2015 Dataset (size 11463×5811) [Perrone et al., 2017]: sampled 10000 rows without replacement.
2. Landmark Dataset (size 71952×2704) (Pereyra/landmark in [Davis and Hu, 2011]): sampled 10000 rows without replacement.




Figure 1: Error rates of *wlr iter*, *svd*, *wsvd*, *rwlra sk* as k' is increased - synthetic datasets.




Figure 2: Errors of *wlra-iter*, *svd*, *wsvd*, *rwlra-sk* as SNR is increased - synthetic datasets

3. Symmetric Stiffness Matrix, Frame Building Dataset (size 1074×1074)(HB/bcsstk08 in [Davis and Hu, 2011]).
4. Blog Feedback DataSet (size 52396×280) [Buza, 2014].

We standardize features of each dataset by removing the mean and scaling to unit variance. We generate weight matrices corresponding to each dataset with following three configurations.

- W_1 : Each element is sampled from $\{1, 0.1, 0.01\}$ with probabilities $\{0.85, 0.1, 0.05\}$.
- W_2 : Each element is sampled from the interval $[0, 1]$ uniformly at random.
- W_3 : A random binary matrix is first chosen by setting each entry to 1 with probability 0.1 and 0 otherwise. Following this, the first 30% columns of first 90% rows are set to 1.

We plot the error with k' in the list $(10, 20, 30, 50, 70)$ and show how the error changes for each algorithm. Similar to synthetic data experiments, we measure the scaled error $(\sum_{ij} W_{ij}(A_{ij} - Z_{ij})^2) / \|A\|_F^2$ where Z is the solution output by each algorithm. We average results over 10 independent runs. Figures 3, 4, 5, 6 show the how the error changes with k' in each dataset.




Figure 3: Errors of *wlra-iter*, *svd*, *wsvd* as k' is increased - NIPS Conference Papers 1987-2015 Dataset.




Figure 4: Errors of *wlra-iter*, *svd*, *wsvd* as k' is increased - Landmark Dataset.

5 Conclusion

We study a natural greedy algorithm for the weighted low rank approximation problem and establish novel additive error guarantees in ℓ_2 and ℓ_p norms for $p > 2$ under a new, realistic, assumption




Figure 5: Errors of *wbra-iter*, *svd*, *wsvd* as k' is increased - Symmetric Stiffness Matrix, Frame Building Dataset.




Figure 6: Errors of *wbra-iter*, *svd*, *wsvd* as k' is increased - Blog Feedback Data Set.

on the target low rank matrix. Our algorithm is easy to implement and works well in practice, compared to natural baselines and previous approaches.

Acknowledgements

Aditya Bhaskara is partially supported by NSF CCF-2008688, an NSF CAREER Award (CCF-2047288) and by a Google Faculty Research Award.

References

- [Adil et al., 2019] Adil, D., Kyng, R., Peng, R., and Sachdeva, S. (2019). Iterative refinement for lp-norm regression. In *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '19, page 1405–1424, USA.
- [Altschuler et al., 2016] Altschuler, J., Bhaskara, A., Fu, G., Mirrokni, V., Rostamizadeh, A., and Zadimoghaddam, M. (2016). Greedy column subset selection: New bounds and distributed algorithms. In *Proceedings of the 33rd International Conference on International Conference on Machine Learning - Volume 48*, ICML'16, page 2539–2548. JMLR.org.
- [Awerbuch and Kleinberg, 2004] Awerbuch, B. and Kleinberg, R. D. (2004). Adaptive routing with end-to-end feedback: Distributed learning and geometric approaches. In *Proceedings of the*

- Thirty-Sixth Annual ACM Symposium on Theory of Computing*, STOC '04, page 45–53, New York, NY, USA. Association for Computing Machinery.
- [Ban et al., 2019a] Ban, F., Bhattachiprolu, V., Bringmann, K., Kolev, P., Lee, E., and Woodruff, D. P. (2019a). A ptas for ℓ_p -low rank approximation. In Chan, T. M., editor, *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2019, San Diego, California, USA, January 6-9, 2019*, pages 747–766. SIAM.
- [Ban et al., 2019b] Ban, F., Woodruff, D., and Zhang, R. (2019b). Regularized weighted low rank approximation. *Advances in Neural Information Processing Systems*, 32:4059–4069.
- [Barak et al., 2012] Barak, B., Brandao, F. G., Harrow, A. W., Kelner, J., Steurer, D., and Zhou, Y. (2012). Hypercontractivity, sum-of-squares proofs, and their applications. *Proceedings of the 44th symposium on Theory of Computing - STOC '12*.
- [Bhaskara and Tai, 2019] Bhaskara, A. and Tai, W. M. (2019). Approximate Guarantees for Dictionary Learning. In Beygelzimer, A. and Hsu, D., editors, *Proceedings of the Thirty-Second Conference on Learning Theory*, volume 99 of *Proceedings of Machine Learning Research*, pages 299–317, Phoenix, USA. PMLR.
- [Bhattachiprolu et al., 2019] Bhattachiprolu, V., Ghosh, M., Guruswami, V., Lee, E., and Tulsiani, M. (2019). Approximability of $p \rightarrow q$ matrix norms: Generalized krivine rounding and hypercontractive hardness. In Chan, T. M., editor, *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2019, San Diego, California, USA, January 6-9, 2019*, pages 1358–1368. SIAM.
- [Bubeck et al., 2018] Bubeck, S., Cohen, M. B., Lee, Y. T., and Li, Y. (2018). An homotopy method for lp regression provably beyond self-concordance and in input-sparsity time. In *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2018, page 1130–1137, New York, NY, USA. Association for Computing Machinery.
- [Buza, 2014] Buza, K. (2014). Feedback prediction for blogs. In *Data analysis, machine learning and knowledge discovery*, pages 145–152. Springer.
- [Candes and Recht, 2008] Candes, E. J. and Recht, B. (2008). Exact matrix completion via convex optimization.
- [Chierichetti et al., 2017] Chierichetti, F., Gollapudi, S., Kumar, R., Lattanzi, S., Panigrahy, R., and Woodruff, D. P. (2017). Algorithms for ℓ_p low-rank approximation. In Precup, D. and Teh, Y. W., editors, *Proceedings of the 34th International Conference on Machine Learning*, volume 70 of *Proceedings of Machine Learning Research*, pages 806–814, International Convention Centre, Sydney, Australia. PMLR.
- [Clarkson, 2010] Clarkson, K. L. (2010). Coresets, sparse greedy approximation, and the frank-wolfe algorithm. *ACM Trans. Algorithms*, 6(4).
- [Clarkson and Woodruff, 2017] Clarkson, K. L. and Woodruff, D. P. (2017). Low-rank approximation and regression in input sparsity time. *J. ACM*, 63(6).

- [Davis and Hu, 2011] Davis, T. A. and Hu, Y. (2011). The university of florida sparse matrix collection. *ACM Transactions on Mathematical Software (TOMS)*, 38(1):1–25.
- [Eriksson and van den Hengel, 2010] Eriksson, A. and van den Hengel, A. (2010). Efficient computation of robust low-rank matrix approximations in the presence of missing data using the l_1 norm. In *2010 IEEE Computer Society Conference on Computer Vision and Pattern Recognition*, pages 771–778.
- [Frieze et al., 2004] Frieze, A., Kannan, R., and Vempala, S. (2004). Fast monte-carlo algorithms for finding low-rank approximations. *Journal of the ACM*, 51(6):1025–1041.
- [Gillis and Glineur, 2011] Gillis, N. and Glineur, F. (2011). Low-rank matrix approximation with weights or missing data is np-hard. *SIAM Journal on Matrix Analysis and Applications*, 32(4):1149–1165.
- [Hardt et al., 2014] Hardt, M., Meka, R., Raghavendra, P., and Weitz, B. (2014). Computational limits for matrix completion. In Balcan, M. F., Feldman, V., and Szepesvári, C., editors, *Proceedings of The 27th Conference on Learning Theory*, volume 35 of *Proceedings of Machine Learning Research*, pages 703–725, Barcelona, Spain. PMLR.
- [Manton et al., 2003] Manton, J. H., Mahony, R., and Yingbo Hua (2003). The geometry of weighted low-rank approximations. *IEEE Transactions on Signal Processing*, 51(2):500–514.
- [Martini et al., 2001] Martini, H., Swanepoel, K. J., and Weiß, G. (2001). The geometry of minkowski spaces — a survey. part i. *Expositiones Mathematicae*, 19(2):97–142.
- [Musco et al., 2020] Musco, C., Musco, C., and Woodruff, D. P. (2020). Simple heuristics yield provable algorithms for masked low-rank approximation.
- [Nesterov, 1998] Nesterov, Y. (1998). Semidefinite relaxation and nonconvex quadratic optimization. *Optimization Methods and Software*, 9(1-3):141–160.
- [Perrone et al., 2017] Perrone, V., Jenkins, P. A., Spano, D., and Teh, Y. W. (2017). Poisson random fields for dynamic feature models. *Journal of Machine Learning Research*, 18.
- [Razenshteyn et al., 2016] Razenshteyn, I., Song, Z., and Woodruff, D. P. (2016). Weighted low rank approximations with provable guarantees. In *Proceedings of the Forty-Eighth Annual ACM Symposium on Theory of Computing*, STOC ’16, page 250–263, New York, NY, USA. Association for Computing Machinery.
- [Song et al., 2017] Song, Z., Woodruff, D. P., and Zhong, P. (2017). Low rank approximation with entrywise l_1 -norm error. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2017, page 688–701, New York, NY, USA. Association for Computing Machinery.
- [Srebro and Jaakkola, 2003] Srebro, N. and Jaakkola, T. (2003). Weighted low-rank approximations. In *Proceedings of the Twentieth International Conference on International Conference on Machine Learning*, ICML’03, page 720–727. AAAI Press.
- [Steinberg, 2005] Steinberg, D. (2005). Computation of matrix norms with applications to robust optimization. *Research thesis, Technion-Israel University of Technology*, 2.

- [Taylor, 1947] Taylor, A. E. (1947). A geometric theorem and its application to biorthogonal systems. *Bulletin of the American Mathematical Society*, 53(6):614–616.
- [Young, 1941] Young, G. (1941). Maximum likelihood estimation and factor analysis. *Psychometrika*, 6(1):49–53.