

Specifying a positive threshold function via extremal points

Vadim Lozin

Mathematics Institute, University of Warwick, UK

V.LOZIN@WARWICK.AC.UK

Igor Razgon

*Department of Computer Science and Information Systems,
Birkbeck University of London, UK*

IGOR@DCS.BBK.AC.UK

Viktor Zamaraev

Elena Zamaraeva

Mathematics Institute, University of Warwick, UK

V.ZAMARAEV@WARWICK.AC.UK

E.ZAMARAEVA@WARWICK.AC.UK

Nikolai Yu. Zolotykh

*Institute of Information Technology, Mathematics and Mechanics,
Lobachevsky State University of Nizhni Novgorod, Russia*

NIKOLAI.ZOLOTYKH@ITMM.UNN.RU

Editors: Steve Hanneke and Lev Reyzin

Abstract

An extremal point of a positive threshold Boolean function f is either a maximal zero or a minimal one. It is known that if f depends on all its variables, then the set of its extremal points completely specifies f within the universe of threshold functions. However, in some cases, f can be specified by a smaller set. The minimum number of points in such a set is the specification number of f . Hu (1965) showed that the specification number of a threshold function of n variables is at least $n + 1$. Anthony et al. (1995) proved that this bound is attained for nested functions and conjectured that for all other threshold functions the specification number is strictly greater than $n + 1$. In the present paper, we resolve this conjecture negatively by exhibiting threshold Boolean functions of n variables, which are non-nested and for which the specification number is $n + 1$. On the other hand, we show that the set of extremal points satisfies the statement of the conjecture, i.e. a positive threshold Boolean function depending on all its n variables has $n + 1$ extremal points if and only if it is nested. To prove this, we reveal an underlying structure of the set of extremal points.

1. Introduction

A Boolean function is called a *threshold function* (also known as *linearly separable* or a *half-space*) if there exists a hyperplane separating true and false points of the function. Threshold functions play fundamental role in the theory of Boolean functions and they appear in a variety of applications such as electrical engineering, artificial neural networks, reliability theory, game theory etc. (see, for example, (Crama and Hammer, 2011)).

We study the problem of teaching threshold functions in the context of on-line learning with a helpful teacher (Goldman and Kearns, 1995). Speaking informally, *teaching* an unknown function f in a given class is the problem of producing its *teaching* (or *specifying*) set, i.e. a set of points in the domain which uniquely specifies f . In the present paper, the universe is the set of threshold functions and a specifying set for f is a subset S of the points

of the Boolean cube such that f is the only threshold function which is consistent with f on S .

It is not difficult to see that in the worst case the specifying set contains all the 2^n points of the Boolean cube. However, in some cases, a threshold function f can be specified by a smaller set, for instance, when f depends on all its variables and is *positive* (or *increasing*), i.e. a function where an increase of a variable cannot lead to a decrease of the function. In this case, f can be specified by the set of its *extremal points*, i.e. its maximal false and minimal true points, of which there are at most $\lfloor \frac{n+1}{2} \rfloor$ (Anthony et al., 1995). Moreover, this description can also be redundant, i.e. sometimes a positive threshold function f can be specified by a proper subset of its extremal points. The minimum cardinality of a teaching set of f , i.e. the minimum number of points needed to specify f , is the *specification number* of f . The maximum specification number over all functions in a class is the *teaching dimension* of the class.

Hu (1965) showed that the specification number of a threshold function with n variables is at least $n + 1$. Anthony et al. (1995) proved that this bound is attained for so-called nested functions by showing that positive nested functions contain precisely $n + 1$ extremal points. They also conjectured that for all other threshold functions with n variables the specification number is strictly greater than $n + 1$.

Our contribution As our first result, we disprove the conjecture of Anthony et al. (1995) by showing that for any $n \geq 4$ there exist threshold functions with n variables which are non-nested and for which the specification number is $n + 1$.

To state our second result, we observe that for positive nested functions the specifying set coincides with the set of extremal points. This is not the case in our counterexamples to the above conjecture. Therefore, our negative resolution of the conjecture leaves open the question on the number of extremal points: is it true that for any positive threshold function different from nested, the number of extremal points is strictly greater than $n + 1$? In this paper, we answer this question positively. Moreover, we prove a slightly more general result dealing with so-called linear read-once functions, which is an extension of nested functions allowing irrelevant variables (see Section 2 for precise definitions). More formally, we prove that a positive threshold function f with $k \geq 0$ relevant variables has exactly $k + 1$ extremal points if and only if f is linear read-once. Our solution is based on revealing an underlying structure of the set of extremal points.

Related work Upper and lower bounds and the average value for the specification number of a threshold Boolean function are obtained in (Anthony et al., 1995).

A number of papers are devoted to the teaching dimension for the class of threshold functions of k -valued logic, i.e. halfspaces defined on the domain $\{0, 1, \dots, k - 1\}^n$. Upper bounds for the teaching dimension are obtained in (Hegedus, 1994; Chirkov and Zolotykh, 2016). A tight lower bound is stated in (Shevchenko and Zolotykh, 1998). The special case $n = 2$ is considered in (Alekseyev et al., 2015; Zamarraeva, 2016, 2017).

The problem of teaching is closely related to the problem of learning (Angluin, 1988; Aizenstein, 1998). Learning threshold functions with membership or/and equivalence queries is studied by Maass and Turan (1994); Hegedus (1994, 1995); Zolotykh and Shevchenko (1997). Special case $n = 2$ is considered in (Bultman and Maass, 1995). Learning threshold

Boolean functions with small weights is investigated in (Abboud et al., 1999; Abasi et al., 2016).

Teaching or/and learning different classes of read-once (or repetition-free) functions are considered in (Angluin et al., 1993; Bshouty et al., 1995; Chistikov, 2011; Chistikov et al., 2014). The importance of linear read-once functions in learning theory is evidenced, in particular, by their connection with special types of decision lists (Rivest, 1987).

Organization of the paper All preliminary information related to the paper can be found in Section 2. The refutation of the conjecture is presented in Section 3. Section 4 contains the results about extremal points of a threshold function. Section 5 concludes the paper with a number of open problems.

2. Preliminaries

Let $B = \{0, 1\}$. For a point $\mathbf{x} \in B^n$ we denote by $(\mathbf{x})_i$ the i -th coordinate of \mathbf{x} , and by $\bar{\mathbf{x}}$ the point in B^n with $(\bar{\mathbf{x}})_i = 1$ if and only if $(\mathbf{x})_i = 0$ for every $i \in [n]$.

Let $f = f(x_1, \dots, x_n)$ be a Boolean function on B^n . For $k \in [n]$ and $\alpha_k \in \{0, 1\}$ we denote by $f|_{x_k=\alpha_k}$ the Boolean function on B^{n-1} defined as follows:

$$f|_{x_k=\alpha_k}(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) = f(x_1, \dots, x_{k-1}, \alpha_k, x_{k+1}, \dots, x_n).$$

For $i_1, \dots, i_k \in [n]$ and $\alpha_1, \dots, \alpha_k \in \{0, 1\}$ we denote by $f|_{x_{i_1}=\alpha_1, \dots, x_{i_k}=\alpha_k}$ the function $(f|_{x_{i_1}=\alpha_1, \dots, x_{i_{k-1}}=\alpha_{k-1}})|_{x_{i_k}=\alpha_k}$. We say that $f|_{x_{i_1}=\alpha_1, \dots, x_{i_k}=\alpha_k}$ is the *restriction* of f to $x_{i_1} = \alpha_1, \dots, x_{i_k} = \alpha_k$. We also say that a Boolean function g is a *restriction* of a Boolean function f if there exist $i_1, \dots, i_k \in [n]$ and $\alpha_1, \dots, \alpha_k \in \{0, 1\}$ such that $g \equiv f|_{x_{i_1}=\alpha_1, \dots, x_{i_k}=\alpha_k}$, i.e., $g(\mathbf{x}) = f|_{x_{i_1}=\alpha_1, \dots, x_{i_k}=\alpha_k}(\mathbf{x})$ for every $\mathbf{x} \in B^{n-k}$.

Definition 1 A variable x_k is called *irrelevant* for f if $f|_{x_k=1} \equiv f|_{x_k=0}$. Otherwise, x_k is called *relevant* for f . If x_k is irrelevant for f we will also say that f does not depend on x_k .

Following the terminology of (Crama and Hammer, 2011), we say that $\mathbf{x} \in B^n$ is a *true point* of f if $f(\mathbf{x}) = 1$ and that $\mathbf{x} \in B^n$ is a *false point* of f if $f(\mathbf{x}) = 0$.

2.1. Positive functions and extremal points

By \preccurlyeq we denote a partial order over the set B^n , induced by inclusion in the power set lattice of the n -set. In other words, $\mathbf{x} \preccurlyeq \mathbf{y}$ if $(\mathbf{x})_i = 1$ implies $(\mathbf{y})_i = 1$. In this case we will say that \mathbf{x} is *below* \mathbf{y} . When $\mathbf{x} \preccurlyeq \mathbf{y}$ and $\mathbf{x} \neq \mathbf{y}$ we will sometimes write $\mathbf{x} \prec \mathbf{y}$.

Definition 2 A Boolean function f is called *positive monotone* (or simply *positive*) if $f(\mathbf{x}) = 1$ and $\mathbf{x} \preccurlyeq \mathbf{y}$ imply $f(\mathbf{y}) = 1$.

For a positive Boolean function f , the set of its false points forms a down-set and the set of its true points forms an up-set of the partially ordered set (B^n, \preccurlyeq) . We denote by

Z^f the set of maximal false points,

U^f the set of minimal true points.

We will refer to a point in Z^f as a *maximal zero of f* and to a point in U^f as a *minimal one of f* . A point will be called an *extremal point of f* if it is either a maximal zero or a minimal one of f . We denote by

$r(f)$ the number of extremal points of f .

2.2. Threshold functions

Definition 3 A Boolean function f on B^n is called a *threshold function* if there exist n weights $w_1, \dots, w_n \in \mathbb{R}$ and a threshold $t \in \mathbb{R}$ such that, for all $(x_1, \dots, x_n) \in B^n$,

$$f(x_1, \dots, x_n) = 0 \iff \sum_{i=1}^n w_i x_i \leq t.$$

The inequality $w_1 x_1 + \dots + w_n x_n \leq t$ is called *threshold inequality* representing function f . It is not hard to see that there are uncountably many different threshold inequalities representing a given threshold function, and if there exists an inequality with non-negative weights, then f is a positive function.

Let $k \in \mathbb{N}, k \geq 2$. A Boolean function f on B^n is *k -summable* if, for some $r \in \{2, \dots, k\}$, there exist r (not necessarily distinct) false points $\mathbf{x}_1, \dots, \mathbf{x}_r$ and r (not necessarily distinct) true points $\mathbf{y}_1, \dots, \mathbf{y}_r$ such that $\sum_{i=1}^r \mathbf{x}_i = \sum_{i=1}^r \mathbf{y}_i$ (where the summation is over \mathbb{R}^n). A function is *asummable* if it is not k -summable for all $k \geq 2$.

Theorem 4 (Elgot, 1961) A Boolean function is a threshold function if and only if it is asummable.

2.3. Linear read-once functions and nested functions

A Boolean function f is called *linear read-once* if it is either a constant function, or it can be represented by a *nested formula* defined recursively as follows:

1. both literals x and \bar{x} are nested formulas;
2. $x \vee t$, $x \wedge t$, $\bar{x} \vee t$, $\bar{x} \wedge t$ are nested formulas, where x is a variable and t is a nested formula that contains neither x , nor \bar{x} .

Eiter et al. (2002) showed that the class of linear read-once functions is precisely the intersection of threshold and read-once functions.

A linear read-once function is called *nested* if it depends on all its variables. For example, the function $(x_1 \vee x_2)x_3x_5$ considered as a function of 5 variables x_1, \dots, x_5 is linear read-once, but not nested, since x_4 is an irrelevant variable. If this function is considered as a function of 4 variables x_1, x_2, x_3, x_5 , then all its variables are relevant and therefore the function is also nested.

It is not difficult to see that a linear read-once function f is positive if and only if a nested formula representing f does not contain negations.

2.4. Specifying sets and specification number

Let \mathcal{F} be a class of Boolean functions of n variables, and let $f \in \mathcal{F}$.

Definition 5 *A set of points $S \subseteq B^n$ is a specifying set for f in \mathcal{F} if the only function in \mathcal{F} consistent with f on S is f itself. In this case we also say that S specifies f in the class \mathcal{F} . The minimal cardinality of specifying set for f in \mathcal{F} is called the specification number of f (in \mathcal{F}) and denoted $\sigma_{\mathcal{F}}(f)$.*

Let \mathcal{H}_n be the class of threshold Boolean functions of n variables. [Hu \(1965\)](#) and later [Anthony et al. \(1995\)](#) showed that the specification number of a threshold function of n variables is at least $n + 1$.

Theorem 6 ([Hu, 1965](#); [Anthony et al., 1995](#)) *For any threshold Boolean function f of n variables $\sigma_{\mathcal{H}_n}(f) \geq n + 1$.*

It was also shown in ([Anthony et al., 1995](#)) that the nested functions attain the lower bound.

Theorem 7 ([Anthony et al., 1995](#)) *For any nested function f of n variables $\sigma_{\mathcal{H}_n}(f) = n + 1$.*

2.5. Essential points

In estimating the specification number of a threshold Boolean function $f \in \mathcal{H}_n$ it is often useful to consider essential points of f defined as follows.

Definition 8 *A point x is essential for f (with respect to class \mathcal{H}_n), if there exists a function $g \in \mathcal{H}_n$ such that $g(x) \neq f(x)$ and $g(y) = f(y)$ for every $y \in B^n$, $y \neq x$.*

Clearly, any specifying set for f must contain all essential points for f . It turns out that the essential points alone are sufficient to specify f in \mathcal{H}_n ([Cover, 1965](#)). Therefore, we have the following well-known result.

Theorem 9 ([Cover, 1965](#)) *The specification number $\sigma_{\mathcal{H}_n}(f)$ of a function $f \in \mathcal{H}_n$ is equal to the number of essential points of f .*

2.6. The number of essential points versus the number of extremal points

It was observed in ([Anthony et al., 1995](#)) that in the study of specification number of threshold functions, one can be restricted to positive functions. To prove Theorem 7, [Anthony et al. \(1995\)](#) first showed that for a positive threshold function f , which depends on all its variables, the set $Z^f \cup U^f$ of extremal points specifies f . Then they proved that for any positive nested function f of n variables $|Z^f \cup U^f| = n + 1$.

In addition to proving Theorem 7, [Anthony et al. \(1995\)](#) also conjectured that nested functions are the only functions with the specification number $n + 1$ in the class \mathcal{H}_n .

Conjecture 10 ([Anthony et al., 1995](#)) *If $f \in \mathcal{H}_n$ has the specification number $n + 1$, then f is nested.*

In the present paper, we disprove Conjecture 10 by demonstrating for every $n \geq 4$ a threshold non-nested function of n variables with the specification number $n + 1$.

On the other hand, we show that the conjecture becomes a true statement if we replace ‘specification number’ by ‘number of extremal points’. In fact, we prove a more general result saying that a positive threshold function f with k relevant variables is linear read-once if and only if it has exactly $k + 1$ extremal points. For this purpose, the following special type of functions appears to be technically useful.

Definition 11 *We say that a Boolean function $f = f(x_1, \dots, x_n)$ is split if there exists $i \in [n]$ such that $f|_{x_i=0} \equiv 0$ or $f|_{x_i=1} \equiv 1$.*

In what follows, we will need the next two observations, which can be easily verified.

Observation 1 *Any positive linear read-once function is split.*

Observation 2 *Any restriction of a linear read-once function is also linear read-once.*

3. Non-nested functions with small specification number

In this section, we disprove Conjecture 10. To this end, we show in the following theorem that the minimum value of the specification number is attained in the class of threshold functions not only by nested functions.

Theorem 12 *For a natural number n , $n \geq 4$ let $f_n = f(x_1, \dots, x_n)$ be a function defined by its DNF*

$$x_1x_2 \vee x_1x_3 \vee \dots \vee x_1x_{n-1} \vee x_2x_3 \dots x_n.$$

Then f_n is positive, not linear read-once, threshold function, depending on all its variables, and the specification number of f_n is $n + 1$.

Proof Clearly, f_n depends on all its variables. Furthermore, f_n is positive, since its DNF contains no negation of a variable. Also, it is easy to verify that f is not split, and therefore by Observation 1 f is not linear read-once.

Now, we claim that the CNF of f_n is

$$(x_1 \vee x_2)(x_1 \vee x_3) \dots (x_1 \vee x_n)(x_2 \vee x_3 \vee \dots \vee x_{n-1}).$$

Indeed, the equivalence of the DNF and CNF can be directly checked by expanding the latter and applying the absorption law:

$$\begin{aligned} & (x_1 \vee x_2)(x_1 \vee x_3) \dots (x_1 \vee x_n)(x_2 \vee x_3 \vee \dots \vee x_{n-1}) \\ &= (x_1 \vee x_2x_3 \dots x_n)(x_2 \vee x_3 \vee \dots \vee x_{n-1}) \\ &= x_1x_2 \vee x_1x_3 \vee \dots \vee x_1x_{n-1} \vee x_2x_3 \dots x_n. \end{aligned}$$

From the DNF and the CNF of f_n we retrieve the minimal ones

$$\begin{aligned} \mathbf{x}_1 &= (1, 1, 0, \dots, 0, 0), \\ \mathbf{x}_2 &= (1, 0, 1, \dots, 0, 0), \\ &\dots \\ \mathbf{x}_{n-2} &= (1, 0, 0, \dots, 1, 0), \\ \mathbf{x}_{n-1} &= (0, 1, 1, \dots, 1, 1) \end{aligned}$$

and maximal zeros of f_n

$$\begin{aligned}\mathbf{y}_1 &= (0, 0, 1, \dots, 1, 1), \\ \mathbf{y}_2 &= (0, 1, 0, \dots, 1, 1), \\ &\dots \\ \mathbf{y}_{n-2} &= (0, 1, 1, \dots, 0, 1), \\ \mathbf{z}_1 &= (0, 1, 1, \dots, 1, 0), \\ \mathbf{z}_2 &= (1, 0, 0, \dots, 0, 1),\end{aligned}$$

respectively (see Theorems 1.26, 1.27 in (Crama and Hammer, 2011)). It is easy to check that all minimal ones $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n-1}$ satisfy the equation

$$(2n - 5)x_1 + 2(x_2 + x_3 + \dots + x_{n-1}) + x_n = 2n - 3,$$

and all maximal zeros $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{n-2}, \mathbf{z}_1, \mathbf{z}_2$ satisfy the inequality

$$(2n - 5)x_1 + 2(x_2 + x_3 + \dots + x_{n-1}) + x_n \leq 2n - 4.$$

Hence the latter is a threshold inequality representing f_n .

Since for a positive threshold function f which depends on all its variables the set of extremal points specifies f , and every essential point of f must belong to each specifying set, we conclude that every essential point of f_n is extremal.

Let us show that the points $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{n-2}$ are not essential for f_n . Suppose to the contrary that there exists a threshold function g_i that differs from f_n only in the point \mathbf{y}_i , $i \in [n-2]$, i.e., $g_i(\mathbf{y}_i) = 1$ and $g_i(\mathbf{x}) = f_n(\mathbf{x})$ for every $\mathbf{x} \neq \mathbf{y}_i$. Then $\mathbf{x}_i + \mathbf{y}_i = \mathbf{z}_1 + \mathbf{z}_2$, and hence g_i is 2-summable. Therefore by Theorem 4 function g_i is not threshold. A contradiction.

The above discussion together with Theorems 6 and 9 imply that all the remaining $n+1$ extremal points $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n-1}, \mathbf{z}_1, \mathbf{z}_2$ are essential, and therefore $\sigma_{\mathcal{H}_n}(f_n) = n+1$. ■

4. Extremal points of a threshold function

The main goal of this section is to prove the following theorem.

Theorem 13 *Let $f = f(x_1, \dots, x_n)$ be a positive threshold function with $k \geq 0$ relevant variables. Then the number of extremal points of f is at least $k+1$. Moreover f has exactly $k+1$ extremal points if and only if f is linear read-once.*

We will prove Theorem 13 by induction on n . The statement is easily verifiable for $n = 1$. Let $n > 1$ and assume that the theorem is true for functions of at most $n-1$ variables. In the rest of the section we prove the statement for n -variable functions. Our strategy consists of three major steps. First, we prove the statement for split functions in Section 4.2. This case includes linear read-once functions. Then, in Section 4.3, we prove the result for non-split functions f which have a variable x_i such that both restrictions $f|_{x_i=0}$ and $f|_{x_i=1}$ are split. Finally, in Section 4.4, we consider the case of non-split functions f , where for every variable x_i of f at least one of the restrictions $f|_{x_i=0}$ and $f|_{x_i=1}$ is non-split. In this case, the proof is based on a structural characterization of the set of extremal points, which is of independent interest and which is presented in Section 4.1.

4.1. The structure of the set of extremal points

We say that a maximal zero (*resp.* minimal one) \mathbf{y} of $f(x_1, \dots, x_n)$ corresponds to a variable x_i if $(\mathbf{y})_i = 0$ (*resp.* $(\mathbf{y})_i = 1$). A pair (\mathbf{a}, \mathbf{b}) of points in B^n is called x_i -extremal for f if

1. \mathbf{a} is a maximal zero of f corresponding to x_i ;
2. \mathbf{b} is a minimal one of f corresponding to x_i ; and
3. $(\mathbf{a})_j \geq (\mathbf{b})_j$ for every $j \in [n] \setminus \{i\}$.

Claim 1 *Let f be a positive function and $i \in [n]$. Then*

1. *for every maximal zero \mathbf{a} of f corresponding to x_i there exists a minimal one \mathbf{b} of f corresponding to x_i such that (\mathbf{a}, \mathbf{b}) is an x_i -extremal pair for f ;*
2. *for every minimal one \mathbf{b} of f corresponding to x_i there exists a maximal zero \mathbf{a} of f corresponding to x_i such that (\mathbf{a}, \mathbf{b}) is an x_i -extremal pair for f .*

Proof We prove the first part of the claim, the second part can be proved similarly. Consider a maximal zero \mathbf{a} of f corresponding to x_i and the vector \mathbf{b}' such that $(\mathbf{a})_j = (\mathbf{b}')_j$ for all $j \neq i$ and $(\mathbf{b}')_i = 1$. Since $\mathbf{a} \prec \mathbf{b}'$ and \mathbf{a} is a maximal zero, we have $f(\mathbf{b}') = 1$. Let \mathbf{b} be a minimal one of f such that $\mathbf{b} \preccurlyeq \mathbf{b}'$. Then $(\mathbf{b})_i = 1$ for otherwise \mathbf{b} would be below \mathbf{a} , which in turn would contradict positivity of f . Now since \mathbf{a} and \mathbf{b}' differ only in coordinate i and $\mathbf{b} \preccurlyeq \mathbf{b}'$, we conclude that $(\mathbf{a})_j \geq (\mathbf{b})_j$ for every $j \in [n] \setminus \{i\}$, and therefore (\mathbf{a}, \mathbf{b}) is an x_i -extremal pair for f . \blacksquare

Let $g = g(y_1, \dots, y_n)$ be a positive function, and let $\{y_{i_1}, \dots, y_{i_k}\}$ be a subset of the relevant variables of g . For every variable y_{i_j} , $j \in [k]$ we fix an y_{i_j} -extremal pair $(\mathbf{a}_{i_j}, \mathbf{b}_{i_j})$. Now we define a graph $H(g, y_{i_1}, \dots, y_{i_k})$ as an undirected graph with vertex set $\{\mathbf{a}_{i_j}, \mathbf{b}_{i_j} \mid j \in [k]\}$ and edge set $\{\{\mathbf{a}_{i_j}, \mathbf{b}_{i_j}\} \mid j \in [k]\}$. We call $H(g, y_{i_1}, \dots, y_{i_k})$ an extremal graph and observe that this graph is not uniquely defined.

Lemma 14 *If g is a threshold function, then $H = H(g, y_{i_1}, \dots, y_{i_k})$ is an acyclic graph.*

Proof It follows from the definitions of an x_i -extremal pair and of an extremal graph that H does not have multiple edges and that H is a bipartite graph with parts $A = \{\mathbf{a}_{i_j} \mid j \in [k]\}$ and $B = \{\mathbf{b}_{i_j} \mid j \in [k]\}$. Suppose to the contrary that H has a cycle of length $2r$, for some $r \in \{2, \dots, k\}$. Let R and Q be the sets of vertices of the cycle belonging to A and B , respectively. For $i \in [n]$ and $\alpha \in \{0, 1\}$ we denote by R_α^i the set of vertices $\mathbf{y} \in R$ with $(\mathbf{y})_i = \alpha$. Similarly, Q_α^i denotes the set of vertices $\mathbf{y} \in Q$ with $(\mathbf{y})_i = \alpha$.

Fix an index $i \in [n]$. By definition of an x_i -extremal pair and of an extremal graph, there is at most one edge between the vertices of Q_1^i and the vertices of R_0^i . Therefore, the number $2|Q_1^i|$ of the edges in the cycle incident to the vertices in Q_1^i is at most one more than the number $2|R_0^i|$ of the edges incident to the vertices in R_0^i . This implies that $|Q_1^i| \leq |R_0^i|$. If this inequality is strict, we modify the set Q by choosing arbitrarily $|R_0^i| - |Q_1^i|$ points in Q_0^i and changing their i -th coordinates from 0 to 1. Since g is positive, the modified points remain true points for g .

Applying this procedure for each $i \in [n]$, we obtain the set R of false points and the set Q of true points both of size r such that $|Q_1^i| = |R_1^i|$ for all i . Therefore, $\sum_{\mathbf{x} \in R} \mathbf{x} = \sum_{\mathbf{y} \in Q} \mathbf{y}$, showing that g is k -summable. Hence, by Theorem 4, g is not threshold, which contradicts the assumption of the lemma. \blacksquare

4.2. Split functions

Lemma 15 *Let $f = f(x_1, \dots, x_n)$ be a positive threshold split function with $k \geq 0$ relevant variables. Then the number of extremal points of f is at least $k + 1$. Moreover f has exactly $k + 1$ extremal points if and only if f is linear read-once.*

Proof The case $k = 0$ is trivial, and therefore we assume that $k \geq 1$.

Let x_i be a variable of f such that $f|_{x_i=0} \equiv \mathbf{0}$ (the case $f|_{x_i=1} \equiv \mathbf{1}$ is similar). Let $f_0 = f|_{x_i=0}$ and $f_1 = f|_{x_i=1}$. Clearly, x_i is a relevant variable of f , otherwise $f \equiv \mathbf{0}$, that is, $k = 0$. Since every relevant variable of f is relevant for at least one of the functions f_0 and f_1 , we conclude that f_1 has $k - 1$ relevant variables.

The equivalence $f_0 \equiv \mathbf{0}$ implies that for every extremal point $(\alpha_1, \dots, \alpha_{i-1}, \alpha_{i+1}, \dots, \alpha_n)$ of f_1 , the corresponding point $(\alpha_1, \dots, \alpha_{i-1}, 1, \alpha_{i+1}, \dots, \alpha_n)$ is extremal for f . For the same reason, there is only one extremal point of f with the i -th coordinate being equal to zero, namely, the point with all coordinates equal to one, except for the i -th coordinate. Hence, $r(f) = r(f_1) + 1$.

1. If f_1 is linear read-once, then f is also linear read-once, since f can be expressed as $x_i \wedge f_1$. By the induction hypothesis $r(f_1) = k$ and therefore $r(f) = k + 1$.
2. If f_1 is not linear read-once, then from Observation 2 we conclude that f is also not linear read-once. By the induction hypothesis $r(f_1) > k$ and therefore $r(f) > k + 1$.

\blacksquare

4.3. Non-split functions with split restrictions

Claim 2 *Let $f = f(x_1, \dots, x_n)$ be a positive threshold non-split function. If there exists $i \in [n]$ such that both $f_0 = f|_{x_i=0}$ and $f_1 = f|_{x_i=1}$ are split, then there exists $s \in [n] \setminus \{i\}$ such that $f_0|_{x_s=0} \equiv \mathbf{0}$ and $f_1|_{x_s=1} \equiv \mathbf{1}$.*

Proof Since f_0 is split, there exists $p \in [n]$ such that $f_0|_{x_p=0} \equiv \mathbf{0}$ or $f_0|_{x_p=1} \equiv \mathbf{1}$. We claim that the latter case is impossible. Indeed, as $f_0|_{x_p=1} = f|_{x_i=0, x_p=1}$, positivity of f and $f_0|_{x_p=1} \equiv \mathbf{1}$ imply $f|_{x_i=1, x_p=1} \equiv \mathbf{1}$, and therefore $f|_{x_p=1} \equiv \mathbf{1}$. This contradicts the assumption that f is non-split. Hence, $f_0|_{x_p=0} \equiv \mathbf{0}$. Similarly, one can show that $f_1|_{x_r=1} \equiv \mathbf{1}$ for some $r \in [n]$. If $p = r$, then we are done.

Assume that $p \neq r$. Let \mathbf{a} be the point in B^n that has exactly two 1's in coordinates i and p . If $f(\mathbf{a}) = 1$, then by positivity $f|_{x_i=1, x_p=1} = f_1|_{x_p=1} \equiv 1$, and the claim follows for $s = p$. Let now \mathbf{b} be a point in B^n that has exactly two 0's in coordinates i and r . If $f(\mathbf{b}) = 0$, then by positivity $f|_{x_i=0, x_r=0} = f_0|_{x_r=0} \equiv 0$, and the claim follows for $s = r$.

Assume now that $f(\mathbf{a}) = 0$ and $f(\mathbf{b}) = 1$. Since $f_{0|x_p=0} \equiv \mathbf{0}$ and $f_{1|x_r=1} \equiv \mathbf{1}$ we conclude that $f(\bar{\mathbf{a}}) = 0$ and $f(\bar{\mathbf{b}}) = 1$. Therefore, $\mathbf{a} + \bar{\mathbf{a}} = \mathbf{b} + \bar{\mathbf{b}}$ and hence by Theorem 4 f is not threshold. This contradiction completes the proof. \blacksquare

Corollary 16

- (a) Variable x_s from Claim 2 is relevant for both functions f_0 and f_1 .
- (b) If a point $\mathbf{a} = (\alpha_1, \dots, \alpha_{i-1}, \alpha_{i+1}, \dots, \alpha_n) \in B^{n-1}$ is an extremal point of f_{α_i} , $\alpha_i \in \{0, 1\}$, then $\mathbf{a}' = (\alpha_1, \dots, \alpha_{i-1}, \alpha_i, \alpha_{i+1}, \dots, \alpha_{n-1}) \in B^n$ is an extremal point of f .

Proof

- (a) Suppose to the contrary that f_0 does not depend on x_s . Then $f_{0|x_s=1} \equiv f_{0|x_s=0} \equiv \mathbf{0}$, and therefore $f_0 = f_{x_s=0} \equiv 0$, which contradicts the assumption that f is non-split. Similarly, one can show that x_s is relevant for f_1 .
- (b) We prove the statement for $\alpha_i = 1$. For $\alpha_i = 0$ the arguments are symmetric. If \mathbf{a} is a maximal zero of f_1 , then \mathbf{a}' is a maximal zero of f . Indeed, for every point $\mathbf{b}' = (\beta_1, \dots, \beta_{i-1}, \beta_i, \beta_{i+1}, \dots, \beta_n) \in B^n$ such that $\mathbf{a}' \prec \mathbf{b}'$ we have $\beta_i = 1$. Hence $\mathbf{a} \prec \mathbf{b} = (\beta_1, \dots, \beta_{i-1}, \beta_{i+1}, \dots, \beta_n)$, and $f_1(\mathbf{b}) = f(\mathbf{b}')$. Therefore $f(\mathbf{b}') = 0$ would imply that \mathbf{a} is not a maximal zero of f_1 . This contradiction shows that \mathbf{a}' is a maximal zero of f .

Let now \mathbf{a} be a minimal one of f_1 . For convenience, without loss of generality, we assume that $s < i$. Suppose to the contrary, that \mathbf{a}' is not a minimal one of f , i.e., there exists a point $\mathbf{b}' = (\beta_1, \dots, \beta_{i-1}, \beta_i, \beta_{i+1}, \dots, \beta_n) \in B^n$ such that $\mathbf{b}' \prec \mathbf{a}'$ and $f(\mathbf{b}') = 1$. Note that if $\beta_i = 1$, then $\mathbf{b} \prec \mathbf{a}$ and $f(\mathbf{b}') = f_1(\mathbf{b})$, where as before, $\mathbf{b} = (\beta_1, \dots, \beta_{i-1}, \beta_{i+1}, \dots, \beta_n)$. Since \mathbf{a} is a minimal one of f_1 , we conclude that $f_1(\mathbf{b}) = f(\mathbf{b}') = 0$, which is a contradiction. Therefore we assume further that $\beta_i = 0$ and distinguish between two cases:

$\beta_s = 0$. In this case

$$f(\mathbf{b}') = (\bar{\beta}_i \wedge f_0(\mathbf{b})) \vee (\beta_i \wedge f_1(\mathbf{b})) = f_0(\mathbf{b}) = 0,$$

where the latter equality follows from $f_{0|x_s=0} \equiv \mathbf{0}$. This is a contradiction to our assumption that $f(\mathbf{b}') = 1$.

$\beta_s = 1$. In this case, $\alpha_s = 1$. Note that the equivalence $f_{1|x_s=1} \equiv \mathbf{1}$ means that function f_1 takes value 1 on every point with s -th coordinate being equal to 1. Together with the minimality of \mathbf{a} this implies that the only non-zero component of \mathbf{a} is α_s . Hence, the only non-zero component of \mathbf{b}' is β_s . Therefore $f(\mathbf{b}') = 1$ and positivity of f imply $f_{|x_s=1} \equiv \mathbf{1}$, which contradicts the assumption that f is non-split. \blacksquare

Lemma 17 *Let $f = f(x_1, \dots, x_n)$ be a positive threshold non-split function with k relevant variables, and there exists $i \in [n]$ such that both $f_0 = f|_{x_i=0}$ and $f_1 = f|_{x_i=1}$ are split. Then the number of extremal points of f is at least $k + 2$.*

Proof Let $s \in [n] \setminus \{i\}$ be an index guaranteed by Claim 2. Let P, P_0 , and P_1 be the sets of relevant variables of f, f_0 , and f_1 , respectively. Since any relevant variable of f is a relevant variable of at least one of the functions f_0, f_1 and, by Corollary 16 (a), x_s is a relevant variable of both of them, we have

$$k = |P| \leq |P_0 \cup P_1| + 1 = |P_0| + |P_1| - |P_0 \cap P_1| + 1 \leq |P_0| + |P_1|.$$

By the induction hypothesis, $r(f_i) \geq |P_i| + 1$, where $i = 0, 1$. Finally, by Corollary 16 (b) the number $r(f)$ of extremal points of f is at least $r(f_0) + r(f_1) \geq |P_0| + |P_1| + 2 \geq k + 2$. \blacksquare

4.4. Non-split functions without split restrictions

Due to Lemmas 15 and 17 it remains to show the bound for a positive threshold non-split function $f = f(x_1, \dots, x_n)$ such that for every $i \in [n]$ at least one of $f_0 = f|_{x_i=0}$ and $f_1 = f|_{x_i=1}$ is non-split.

Assume without loss of generality that x_n is a relevant variable of f , and let $f_0 = f|_{x_n=0}$ and $f_1 = f|_{x_n=1}$. We assume that f_0 is non-split and prove that f has at least $k + 2$ extremal points, where k is the number of relevant variables of f . The case when f_0 is split, but f_1 is non-split is proved similarly. Let us denote the number of relevant variables of f_0 by m . Clearly, $1 \leq m \leq k - 1$. Exactly $k - 1 - m$ of k relevant variables of f became irrelevant for the function f_0 . Note that these $k - 1 - m$ variables are necessarily relevant for the function f_1 . By the induction hypothesis, the number $r(f_0)$ of extremal points of f_0 is at least $m + 2$.

We introduce the following notation:

C_0 – the set of maximal zeros of f corresponding to x_n ;

P_0 – the set of all other maximal zeros of f , i.e., $P_0 = Z^f \setminus C_0$;

C_1 – the set of minimal ones of f corresponding to x_n ;

P_1 – the set of all other minimal ones of f , i.e., $P_1 = U^f \setminus C_1$.

For a set $A \subseteq B^n$ we will denote by A^* the restriction of A into the first $n - 1$ coordinates, i.e., $A^* = \{(\alpha_1, \dots, \alpha_{n-1}) \mid (\alpha_1, \dots, \alpha_{n-1}, \alpha_n) \in A \text{ for some } \alpha_n \in \{0, 1\}\}$.

By definition, the number of extremal points of f is

$$r(f) = |C_0| + |P_1| + |C_1| + |P_0| = |C_0^*| + |P_1^*| + |C_1^*| + |P_0^*|. \quad (1)$$

We want to express $r(f)$ in terms of the number of extremal points of f_0 and f_1 . For this we need several observations. First, for every extremal point $(\alpha_1, \dots, \alpha_{n-1}, \alpha_n)$ for f the point $(\alpha_1, \dots, \alpha_{n-1})$ is extremal for f_{α_n} . Furthermore, we have the following straightforward claim.

Claim 3 *P_1^* is the set of minimal ones of f_0 and P_0^* is the set of maximal zeros of f_1 .*

In contrast to minimal ones of f_0 , the set of maximal zeros of f_0 in addition to the points in C_0^* may contain extra points, which we denote by N_0^* . In other words, $Z^{f_0} = C_0^* \cup N_0^*$. Similarly, besides C_1^* , the set of minimal ones of f_1 may contain additional points, which we denote by N_1^* . That is, $U^{f_0} = C_1^* \cup N_1^*$.

Claim 4 *The set N_0^* is a subset of the set P_0^* of maximal zeros of f_1 . The set N_1^* is a subset of the set P_1^* of minimal ones of f_0 .*

Proof We will prove the first part of the statement, the second one is proved similarly. Suppose to the contrary that there exists a point $\mathbf{a} = (\alpha_1, \dots, \alpha_{n-1}) \in N_0^* \setminus P_0^*$, which is a maximal zero for f_0 , but is not a maximal zero for f_1 . Notice that $f_1(\mathbf{a}) = 0$, as otherwise $(\alpha_1, \dots, \alpha_{n-1}, 0)$ would be a maximal zero for f , which is not the case, since $\mathbf{a} \notin C_0^*$. Since \mathbf{a} is not a maximal zero for f_1 , there exists a maximal zero $\mathbf{b} \in B^{n-1}$ for f_1 such that $\mathbf{a} \prec \mathbf{b}$. But then we have $f_0(\mathbf{b}) = 1$ and $f_1(\mathbf{b}) = 0$, which contradicts positivity of function f . ■

From Claim 3 we have $r(f_0) = |Z^{f_0} \cup U^{f_0}| = |C_0^*| + |N_0^*| + |P_1^*|$, which together with (1) and Claim 4 imply

$$\begin{aligned} r(f) &= |C_0^*| + |P_1^*| + |C_1^*| + |P_0^*| = |C_0^*| + |P_1^*| + |C_1^*| + |N_0^*| + |P_0^* \setminus N_0^*| \\ &= r(f_0) + |C_1^*| + |P_0^* \setminus N_0^*|. \end{aligned} \quad (2)$$

Using the induction hypothesis we conclude that $r(f) \geq m + 2 + |C_1^*| + |P_0^* \setminus N_0^*|$. To derive the desired bound $r(f) \geq k + 2$, in the rest of this section we show that $C_1^* \cup P_0^* \setminus N_0^*$ contains at least $k - m$ points.

Claim 5 *Let x_i , $i \in [n-1]$, be a relevant variable for f_1 , but irrelevant for f_0 . Then there exists an x_i -extremal pair (\mathbf{a}, \mathbf{b}) for f_1 such that $\mathbf{a} \in P_0^* \setminus N_0^*$ and $\mathbf{b} \in C_1^*$.*

Proof First, let us show that an x_i -extremal pair always exists. Since x_i is relevant for f_1 , there exists a pair of points \mathbf{x} and \mathbf{y} , which differ only in the i -th coordinate and $f_1(\mathbf{x}) \neq f_1(\mathbf{y})$. Without loss of generality, let $(\mathbf{x})_i = 0$ and $(\mathbf{y})_i = 1$. Then by positivity, $f_1(\mathbf{x}) = 0$ and $f_1(\mathbf{y}) = 1$. Let \mathbf{x}' be any maximal zero of f_1 such that $\mathbf{x} \preccurlyeq \mathbf{x}'$. Then obviously \mathbf{x}' is a maximal zero corresponding to x_i and the existence of an x_i -extremal pair for f_1 follows from Claim 1.

We claim that $(\mathbf{x})_i = 1$ for every $\mathbf{x} \in N_0^*$. Indeed, if $(\mathbf{x})_i = 0$ for a maximal zero $\mathbf{x} \in N_0^*$, then changing in \mathbf{x} the i -th coordinate from 0 to 1 we would obtain the point \mathbf{x}' with $f_0(\mathbf{x}') = 1 \neq f_0(\mathbf{x})$, which would contradict the assumption that x_i is irrelevant for f_0 . Similarly, one can show that $(\mathbf{y})_i = 0$ for every $\mathbf{y} \in P_1^*$.

The above observations together with Claim 4 imply that every maximal zero for f_1 corresponding to x_i belongs to $P_0^* \setminus N_0^*$ and every minimal one for f_1 corresponding to x_i belongs to C_1^* . Hence the claim. ■

Recall that there are exactly $s = k - 1 - m$ variables that are relevant for f_1 and irrelevant for f_0 . We denote these variables by x_{i_1}, \dots, x_{i_s} . Let H be an extremal graph $H(f_1, x_{i_1}, \dots, x_{i_s})$ defined in such a way that all its vertices belong to $C_1^* \cup P_0^* \setminus N_0^*$. Such a graph exists by Claim 5. By Lemma 14 the graph H is acyclic, and hence it has at least $s + 1$ vertices. Therefore, the set $C_1^* \cup P_0^* \setminus N_0^*$ has at least $s + 1 = k - m$ points. This conclusion establishes the main result of this section.

Lemma 18 *Let $f = f(x_1, \dots, x_n)$ be a positive threshold non-split function with k relevant variables, and for every $i \in [n]$ at least one of the restrictions $f_0 = f|_{x_i=0}$ and $f_1 = f|_{x_i=1}$ is non-split. Then the number of extremal points of f is at least $k + 2$.*

5. Conclusion and open problems

In this paper we studied the cardinality and structure of two sets related to teaching positive threshold Boolean functions: the specifying set and the set of their extremal points.

First, we showed the existence of positive threshold Boolean functions of n variables, which are not linear read-once and for which the *specification number* is at its lowest bound, $n + 1$ (Theorem 12). An important open problem is to describe the set of all such functions.

Second, we completely described the set of all positive threshold Boolean functions of n relevant variables, for which the number of *extremal points* is at its lowest bound, $n + 1$. This is precisely the set of all positive linear read-once functions (Theorem 13). It would be interesting to find out whether this result is valid for all positive functions, not necessarily threshold. In other words, is it true that a positive Boolean function of n relevant variables has $n + 1$ extremal points if and only if it is linear read-once?

Finally, we ask whether the acyclic structure of the set of extremal points of a positive threshold function f can be helpful in determining the specification number of f .

Acknowledgment This work was supported by the Russian Science Foundation Grant No. 17-11-01336.

References

Elias Abboud, Nader Agha, Nader H. Bshouty, Nizar Radwan, and Fathi Saleh. Learning threshold functions with small weights using membership queries. In *Proceedings of the Twelfth Annual Conference on Computational Learning Theory*, pages 318-322, ACM, 1999.

Howard Aizenstein, Tibor Hegedüs, Lisa Hellerstein, and Leonard Pitt. Complexity theoretic hardness results for query learning. *Computational Complexity*, 7(1):19-53, 1998.

Max A. Alekseyev, Marina G. Basova, and Nikolai Yu. Zolotykh. On the minimal teaching sets of two-dimensional threshold functions. *SIAM Journal on Discrete Mathematics*, 29(1):157-165, 2015.

Dana Angluin. Queries and concept learning. *Machine learning*, 2(1):319-342, 1988.

Dana Angluin, Lisa Hellerstein, and Marek Karpinski. Learning read-once formulas with queries. *Journal of the ACM*, 40(1):185-210, 1993.

Martin Anthony, Graham Brightwell, and John Shawe-Taylor. On specifying Boolean functions by labelled examples. *Discrete Applied Mathematics*, 61(1):1-25, 1995.

Hasan Abasi, Ali Z. Abdi, and Nader H. Bshouty. Learning Boolean halfspaces with small weights from membership queries. *Theoretical Computer Science*, 650(18):45-56, 2016.

Nader H. Bshouty, Thomas R. Hancock, and Lisa Hellerstein. Learning arithmetic read-once formulas. *SIAM Journal on Computing*, 24(4):706-735, 1995.

William J. Bultman, and Wolfgang Maass. Fast identification of geometric objects with membership queries. *Information and Computation*, 118(1):48-64, 1995.

Aleksandr Yu. Chirkov, and Nikolai Yu. Zolotykh. On the number of irreducible points in polyhedra. *Graphs and Combinatorics*, 32(5):1789-1803, 2016.

D. V. Chistikov. Testing monotone read-once functions. In *International Workshop on Combinatorial Algorithms*, pages 121-134, Springer, Berlin, Heidelberg, 2011.

Dmitry Chistikov, Valentina Fedorova, and Andrey Voronenko. Certificates of non-membership for classes of read-once functions. *Fundamenta Informaticae*, 132(1):63-77, 2014.

Thomas M. Cover. Geometrical and statistical properties of systems of linear inequalities with applications in pattern recognition. *IEEE Transactions on Electronic Computers*, EC-14(3):326-334, 1965.

Yves Crama, and Peter L. Hammer. *Boolean functions: Theory, algorithms, and applications*. Cambridge University Press, 2011.

Thomas Eiter, Toshihide Ibaraki, and Kazuhisa Makino. Decision lists and related Boolean functions. *Theoretical Computer Science*, 270(1):493-524, 2002.

Calvin C. Elgot. Truth functions realizable by single threshold organs. In *Proceedings of the Second Annual Symposium on Switching Circuit Theory and Logical Design*, pages 225-245, SWCT, 1961.

Sally A. Goldman, and Michael J. Kearns. On the complexity of teaching. *Journal of Computer and System Sciences*, 50(1):20-31, 1995.

Tibor Hegedüs: Geometrical concept learning and convex polytopes. In *Proceedings of the 7th Annual ACM Conference on Computational Learning Theory*, pages 228-236, ACM Press, New York, 1994.

Tibor Hegedüs: Generalized teaching dimensions and the query complexity of learning. In *Proceedings of the 8th Annual ACM Conference on Computational Learning Theory*, pages 108-117, ACM Press, New York, 1995.

Sze-Tsen Hu. *Threshold Logic*. University of California Press, Berkeley, 1965.

Wolfgang Maass, and György Turán. How fast can a threshold gate learn? In *Computational Learning Theory and Natural Learning Systems: Constraints and Prospects*, S. Hanson, G. Drastal, R. Rivest, editors, pages 381-414, MIT Press, 1994.

Ronald L. Rivest. Learning decision lists. *Machine Learning*, 2(3):229-246, 1987.

V. N. Shevchenko, N. Yu. Zolotykh. On complexity of deciphering threshold functions of k -valued logic. *Russian Mathematical Doklady*, 362(5):606-608, 1998.

Elena Zamaraeva. On teaching sets of k -threshold functions. *Information and Computation*, 251:301-313, 2016.

E. M. Zamaraeva. On teaching sets for 2-threshold functions of two variables. *Journal of Applied and Industrial Mathematics*, 11(1):130-144, 2017.

N. Yu. Zolotykh, and V. N. Shevchenko. Deciphering threshold functions of k -valued logic. in *Operations Research and Discrete Analysis*, pages 321-326, Kluwer Ac. Publ., Netherlands, 1997.